



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
 Orissa University of Agriculture & Technology,
 Bhubaneswar



Agromet Advisory Bulletin

Date : 12-12-2023

Weather Forecast of District BALANGIR(Odisha) Issued On : 2023-12-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-12-13	2023-12-14	2023-12-15	2023-12-16	2023-12-17
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	27.0	27.0	27.0	28.0	27.0
Tmin(°C)	13.0	11.0	11.0	11.0	12.0
RH-I(%)	62	61	54	51	61
RH-II(%)	32	31	29	24	30
Wind Speed(kmph)	3	4	6	5	5
Wind Direction(Degree)	75	63	65	63	135
Cloud Cover(Octa)	3	3	1	0	0

Weather Summary/Alert:

As per IMD, no rainfall is forecast for upcoming 5 days in the district. The day temperature will be 27-28 °C and night temperature will be 11-13 °C for upcoming 5 days. Maximum and minimum relative humidity will be 51-62 % and 24-32 % respectively. The wind speed will remain within 3 to 6 kmph for upcoming 5 days. There will be mainly clear to partly cloudy sky for next 5 days.

General Advisory:

Keep the harvested product in safe place, carry harvested produce to covered threshing floor. • Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature. • To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cauliflower, Cabbage, Chilly & tomato spray 0.5 g Emamectin benzoate or 0.4 ml Spinosad per liter of water.

SMS Advisory:

Regular surveillance work should be taken up in all crops in order to know the incidence of various disease pests and initiate the control strategies.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	<input type="checkbox"/> Go for threshing of the harvested paddy. <input type="checkbox"/> After threshing, paddy grains need to be sun-dried to 14% moisture for consumption purpose and for seed purpose it should be dried to 12% moisture for better self-life. <input type="checkbox"/> If 'Rodent/Rats' problem is noticed, locate the rodent burrow and surrounding areas. Place Aluminium Phosphide 6% tablet @ One tablet (12 gm) per burrow and seal the burrow with mud which will kill the rodents.

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	There is a chance of powdery mildew disease in Black gram and Green gram. To control this disease, spray Sulphur 80% WP @800 g/acre or Hexaconazole 5% E.C @ 300 ml/acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
CAULIFLOWER	To manage diamond back moth install 6-8 nos of light traps per acre. During primary stage of pest infestation spray neembased pesticide (Azadirachtin) 300 PPM @ 1-litre/acre by mixing in 200-litre of water. To manage this pest chemically spray Tolfenpyrad 15 % EC @ 400-ml/acre or Flubendiamide 20 % WG @ 20-gram/acre or Emamectin Benzoate 5 % SG @ @80-gram/acre by mixing in 200-litre of water.
POTATO	There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Goat kids suffer from Enterotoxemia if graze dew wet grass in winter morning. By this large number of parasites enter into Kid stomach leading Enterotoxemia causing sudden death. To prevent this kids should not be allowed to graze during early morning in winter season. Kids should be allowed to graze after dew gets dried off.