

Gramin Krishi Mausam Sewa India Meteorological Department Odisha University of Agriculture and Technology Ranital, Bhadrak, - 756111

WeekNo.49

No.-98 (Eng.Bulletin) Dt.-08.12.2023

District-Bhadrak (North-East Coastal Plain Zone)

Weather Condition During Last Week at Bhadrak (02.12.2023 –08.12.2023)

Rainfall (mm)	5.3
Maximum temperature (⁰ C)	23.0-30.0
Minimum temperature (⁰ C)	20.0-23.0

<u>Forecast (Upto 13.12.2023)</u> <u>Given by Met.Centre,IMD,Bhubaneswar</u>

DISTRICT: Bhadrak – As per IMD forecast, the district is likely to receive Very Light rainfall on first day and dry weather for last 4 days with Mainly clear to Generally Cloudy sky. The wind speed is likely to remain within 08-10 km/h. The maximum temperature and minimum temperature will likely to remain 25-29°C and 16-18°C to respectively for the next five days.

	DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
		09-12-2023	2	25	18	5	80	58	10	36
		10-12-2023	0	27	17	3	72	52	8	22
	BHADRAK	11-12-2023	0	28	16	2	69	42	8	150
		12-12-2023	0	29	16	2	49	31	10	56
		13-12-2023	0	29	16	3	50	26	8	90

For further information, contact the Met.Centre,Aerodrom Area,IMD,Bhubaneswar,Tel.#0674-2596116.

Agromet Advisory

	✤ General Advisory-
Kharif Rice	★ As per IMD there may be chances of Very light rain on first day and dry weather for last 4 days,
	so farmers are advised to go for sowing of rabi crops and winter vegetables.
	• Harvest the matured paddy and keep in safer place.
	Make the threshing floor and storage ground clean and disinfectant the same.
	Clean the storage Go down properly and spray Lambda Cyahalothrin <u>4.9CS@ 1.5ml/li</u> . of water to
	minimize the chance of future infestation of storage pest.
	✤ Grains must be dried under sun for 3-4 days to 14% moisture for consumption and 12% moisture for seed purpose.
Rapeseed and	* Taking the advantage of store soil moisture, land preparation and sowing of the crop should be
Mustard	completed soon to get optimum yield.
	Suitable varieties: Toria- Anuradha, Parbati, Sushree
	Indian mustard- NRCHB 101, Mahyco bold, JD 6
	Seed rate- 8-10 Kg/ha, Spacing- 30 cm X 10 cm
	 Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha
	Mustard@80:40:40 N-P2O5-K2O Kg/ha
	50%N and all P2O5 and K2O fertilizers should be applied as basal.
	♦ Application of Bentonite Sulphur @30kg/ha for increasing the oil content .Otherwise select Grommer (20:20:0:12) and MOD as a source of N D K & S
	(20:20:0:13) and MOP as a source of N,P,K & S.
Vegetables	To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake. And spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.
	 In case of solanaceous vegetable avoid excess irrigation to prevent the occurrence and spreading of wilting disease. If wilting is noticed uproot the plants, expose the soil to sunlight and spray Tebuconazole @200ml along with Plantomycin @ 200 g/acre.

	Provide bamboo or wooden stake for better fruit set and growth of Tomato plants.
	* For minimizing the infestation of sucking insect pests like white fly, leaf hopper, thirps and aphid in
	vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water.
	To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly &
	tomato apply 0.5 g Emamectin benzoate or 0.4 ml Spinosad per liter of water.
	✤ Legume Vegetables- Cowpea, Beans, Runner bean
	To protect the crop from flea beetle and other foliage feeder spray Neem oil 1500 PPM @ 3ml/li. of
	water or spinosad 0.4 ml/lit of water recommended.
	• For Minimizing the spread of viral diseases like Cowpea Mosaic virus, Install yellow sticky trap, spray
	Neem based formulation 1500 ppm @ 3 ml/ lit of H 2O and need based application of Acetamiprid @
	0.4g/lit of water recommended.
Cole Crops	To prevent collar rot of cole crops Spraying of (Metalxyl + Mancozeb) or (Cymaxonil+Mancozeb) @
	2g /lit of water recommended.
<mark>Fruit</mark>	◆ Papaya- Install Yellow Sticky trap to trap the vector (Whitefly) of Papaya Leaf Curl Virus. Alternate
	spraying of Neem oil formulation 1500ppm @3ml/lit of water and flonicamid 50WG @0.3g/litre to
	control whitefly.
	✤ Avoid Growing of crops like – Okra, Cowpea in the near vicinity of papaya crops.
	✤ Coconut-
	* Monitor the activity and infestation level of Red Palm Weevil and Rhinocerous Beetle. Spraying of
	Lambda Cyahalothrin @1.25ml/lit. of water in the manure pits adjoining to coconut plantation
	recommended.
	* To reduce the incidence of Leaf blight in newly planted coconut saplings spraying of Meatalxyl+
	Mancozeb@2 g/ litre of water is advised.
Pisciculture	★ As the winter is in the starting phase, there is a possibility of "EUS" disease in fish. Farmers are
	advised to apply 'Cifax' @ 400 ml mixed with 40 litre of water in one acre pond area as a preventive
	measure.
	✤ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal	Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may
Science	be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of
	the animals.
	◆ 10% extra dry matter should be provided over and above the total requirement to fulfill the extra
	energy requirement in winter

Mrs. Meera Mahanty Jr. Agrometeorologist AMFU, RRTTS, Ranital Dr. Umasankar Nayak ADR & NODAL OFFICER AMFU, RRTTS, Ranital

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi &Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download:(Android:<u>https://play.google.com/store/apps/details?id=com.aas.meghdoot)</u>,(iOS:<u>https://apps.apple.com/in/</u> app/meghdoot/id1474048155)