

Gramin Krishi Mausam Sewa India Meteorological Department Odisha University of Agriculture and Technology Ranital, Bhadrak, - 756111

WeekNo.52

No.-103 (Eng.Bulletin) Dt.-26.12.2023

District—Bhadrak (North-East Coastal Plain Zone)

Weather Condition During Last Week at Bhadrak (20.12.2023 –26.12.2023)

Rainfall (mm)	0.0			
Maximum temperature (°C)	26.0-28.0			
Minimum temperature (⁰ C)	13.0-17.0			

Forecast (Upto 31.12.2023) Given by Met.Centre,IMD,Bhubaneswar

DISTRICT: Bhadrak—As per IMD forecast, the district is likely to receive **Dry weather for 5 days** with Clear sky. The wind speed is likely to remain within 07-09 km/h. The maximum temperature and minimum temperature will likely to remain 28-29°C and 16-17°C to respectively for the next five days.

	o week to the trape the test test test to the test test test to the test test test test test test test								
DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
	27-12-2023	0	29	17	0	75	39	7	288
	28-12-2023	0	28	17	0	71	35	7	120
BHADRAK	29-12-2023	0	28	16	0	67	35	9	68
	30-12-2023	0	28	16	0	61	35	9	112
	31-12-2023	0	28	16	0	62	33	8	105

For further information, contact the Met.Centre, Aerodrom Area, IMD, Bhubaneswar, Tel. #0674-2596116.

Agromet Advisory

		Agromet Advisory
	G	eneral Advisory-
	*	Farmers are advised not to burn the residue of Kharif crops in the fields because it not only pollute the environment but also cause severe health problems.
	*	To prevent damage from fog,low temperature and moisture stress ,provide light irrigation to standing crops & vegetables during morning hours.
Kharif Rice	*	Rice grains must be dried under sun for 3-4 days to 14% moisture for consumption and 12% moisture for seed purpose.
v	*	• •
	*	In the shallow low land areas sowing of local blackgram as paira crop should be completed.
Blackgram	*	Sow the seeds after treating with (Carboxin+Thiram)@2gm/kg of seed.
	*	Treat the seed after 7 days of above insecticide with specific Rhizobium culture before sowing. Mix 400g culture with 3g Ammonium molybdate & 800 ml water and apply with 10 kg of seed.
Rapeseed and	*	Taking the advantage of store soil moisture, land preparation and sowing of the crop should be
Mustard		completed soon to get optimum yield.
riustai a	*	Suitable varieties: Toria- Sushree
		Indian mustard- NRCHB 101, JD 6, Sampoorna
	*	8 4,41
	*	Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha
		Mustard@80:40:40 N-P2O5-K2O Kg/ha
	.•.	50%N and all P2O5 and K2O fertilizers should be applied as basal.
	*	Application of Bentonite Sulphur @30kg/ha for increasing the oil content .Otherwise select Grommer (20:20:0:13) and MOP as a source of N,P,K & S.
	*	Thinning should be carried out before top dressing to maintain optimum plant population.

Vegetables	To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake. And
	spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.
	❖ In case of solanaceous vegetable avoid excess irrigation to prevent the occurrence and spreading of wilting disease. If wilting is noticed uproot the plants, expose the soil to sunlight and spray Tebuconazole @200ml along with Plantomycin @ 200 g/acre.
	❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants.
	For minimizing the infestation of sucking insect pests like white fly, leaf hopper, thirps and aphid in
	vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water.
	To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly &
	tomato apply 0.5 g Emamectin benzoate or 0.4 ml Spinosad per liter of water.
	 ❖ Legume Vegetables- Cowpea, Beans, Runner bean
	To protect the crop from flea beetle and other foliage feeder spray Neem oil 1500 PPM @ 3ml/li. of water or spinosad 0.4 ml/lit of water recommended.
	❖ For Minimizing the spread of viral diseases like Cowpea Mosaic virus, Install yellow sticky trap, spray
	Neem based formulation 1500 ppm @ 3 ml/ lit of H 2O and need based application of Acetamiprid @
	0.4g/lit of water recommended.
Cole Crops	❖ To prevent collar rot of cole crops Spraying of (Metalxyl + Mancozeb) or (Cymaxonil+Mancozeb) @ 2g /lit of water recommended.
Fruit	❖ Papaya-Install Yellow Sticky trap to trap the vector (Whitefly) of Papaya Leaf Curl Virus. Alternate
	spraying of Neem oil formulation 1500ppm @3ml/lit of water and flonicamid 50WG @0.3g/litre to control whitefly.
	❖ Coconut-
	To reduce the incidence of Leaf blight in newly planted coconut saplings spraying of Meatalxyl+ Mancozeb@2 g/ litre of water is advised.
Pisciculture	❖ As the winter is in the starting phase, there is a possibility of "EUS" disease in fish. Farmers are
	advised to apply 'Cifax' @ 400 ml mixed with 40 litre of water in one acre pond area as a preventive
	measure.
	❖ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal	➤ If the animals have not yet been vaccinated against FMD, Haemorrhagic Septicaemia, Black Quarter,
Science	Enterotoxemia etc., these must be done.
	Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may
	be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of
	the animals.

Mrs. Meera Mahanty Jr. Agrometeorologist AMFU, RRTTS, Ranital Dr. Umasankar Nayak ADR & NODAL OFFICER AMFU, RRTTS, Ranital

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

 $\label{lem:combined} \textbf{Download:} (And \textbf{roid:} \underline{\textbf{https://play.google.com/store/apps/details?id=com.aas.meghdoot)}, (iOS:\underline{\textbf{https://apps.apple.com/in/app/meghdoot/id1474048155}})$