



Gramin Krishi Mausam Sewa
India Meteorological Department
Orissa University of Agriculture and Technology,
Regional Research & Technology Transfer Sub Station,
Kirci, Sundergarh 770073



Dr. Sajeeb Kumar Biswasi,
Senior Scientist & Nodal Officer, GKMS

No-01(Eng.Bulletin)
Dt -02.01.2024

Week No -01

District –Deogarh (North Western Plateau Agroclimatic Zone)

The mean maximum daily temperature was 27.6⁰C & mean minimum daily temperature was 15.2⁰C in Deogarh district during last week. The District received 0.0mm rainfall during the last week. The kharif Paddy is at maturity to harvesting stage. Early sown non-Paddy crops like Pulses, Oilseeds crop are at harvesting stage. The Vegetables are at harvesting stage. The overall crop condition is normal.

Forecast (Up to 07.01.2024)

Given by Met.Centre, IMD, Bhubaneswar

DISTRICT: DEOGARH As per IMD Forecast, the district is may likely receive verylight rainfall for the next five days. The sky condition may be clear sky for the next five days .The wind speed is likely to remain within 5-6km/h. The Maximum Temperature and Minimum Temperature will likely to remain between 22-24⁰C and 15-17⁰C respectively for the next five days.

DISTRICT	03/12/2024	04/12/2024	05/01/2024	06/01/2024	07/01/2024
Date	03/12/2024	04/12/2024	05/01/2024	06/01/2024	07/01/2024
Rainfall (mm)	0	0	0	1	3
T.MAX(°C)	24	24	23	22	23
T.MIN (°C)	15	16	17	17	17
Cloud Cover	0	0	0	0	0
Rh Max (%)	69	72	81	80	81
Rh Min (%)	27	29	27	31	32
Wind speed (kmph)	5	5	5	6	5
Wind Direction (deg)	72	116	135	108	117

For further information,contact the Met. Centre,Aerodrom Area, IMD, Bhubaneswar, Tel.# 0674-2596116

Agromet Advisory

DISTRICT:SUNDERGARH:-

Paira Crop : Sow the seeds of pulses in paddy field utilizing the residual soil moisture before 15 days of harvestingthe paddy. The pulses grown as of paira crop to be sprayed with 2% DAP solution for crop vigour.

Rabi Paddy : Prepare the rabi paddy nursery. Chose medium duration variety. Prepare beds of 1 m width and length according to availability and 10 decimil for one acre land. 40 baskets of FYM and 12 kg of SSP and 2 kg MOP is required for nursery beds. Sow the presoaked seeds on the bed.

Wheat:

The Crown Root Initiation at 15-21 days after sowing is critical for irrigation. Provide irrigation at this stage. Apply 15-20 kg N per acre at 35 days after sowing.

Groundnut Prepare the land for groundnut cultivation. Chose the sandy loam soil having irrigation facility. The winter groundnut at 35-40 days after sowing do not disturb the soil due to peg formation stage.

Sweetpotatoes :

When the sweetpotatoes are 25 to 30 days old, weed the crop and fertilize it with 37.5 kg of potassium per acre and 37.5 kg of potash fertilizer.

Irrigate by looking at the soil moisture through the fertilizer

Sunflower :

Reduce thrips infestation by planting crops such as corn, carrots, and millet around sun-sensitive crops.

For better yield, add manure and root fertilizer to the soil a week before sowing

Keep the field weed free for the first 45 days

Do not apply pesticides to sunflower crops at all

Spray 400 ml of dimethoate per acre in 200 liters of water for sucking insects such as aphids.

Potato: Without any delay complete sowing of Potato. Seed rate is 6-8q/ac. Good varieties are Kufri Pokhraj, Kufri Lalima, Kufri chipsona. The seed weight should be 25-35 g per piece and having 2-3 eyes.

Animal: Keep clean and dry the animal shed. Arrange smoke of neem leaves near to animal shed for preventive measures of diseases. Provide need based animal vaccination in consultation with line department. Animals should be given the wheat grains, jaggery and the mixture of mineral with white salt at the daily basis all these may be provided @ 05-15% in the daily ration basis. Due to forecast of decreasing temperature, cover the animal shed specially in nights.

Poultry :

For more eggs and meat, keep Banraja, Grampriya breeds of chickens.

Vaccinate chickens regularly against distemper and vernal distemper as advised by a veterinarian.

For any problem related to agriculture sector, please dial the toll free number 1551.

For better agricultural benefits use, **OUAT KALINGA** Products (Seedlings, Saapplings and QPM, Tissue culture plants, Fruits and vegetables, value added products, Fingerlings/Yearlings, Poultry chicks, Breeds, Mushroom and spwn, Biofertilizer, Biopesticides, Vermi & Vermicompost and Farm implements)

IMD weather forecast & Agrometeorological Advisory of Odisha location is now available on Meghdoot mobile app in English & Odia Language.

Download: (Andriod: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>)

ବି : ଦ୍ରଷ୍ଟବ୍ୟ :ଓଡ଼ିଆ ଶାଖା ଚିପଲିମା ଓ ଗନ୍ଧାରିପାଲି ବିହନ ଗବେଷଣା ଫାର୍ମ ଠାରେ ସ୍ୱର୍ଣ୍ଣ ଧାନ ବିହନ ୨୦ କିଗ୍ରା ବସ୍ତା ପିଛା ଟ. ୭୧୨ ୪୦ . ପଇସା ଚାଷୀ ନିମିତ୍ତ ବିକ୍ରି ହେଉଅଛି ।

Nodal Officer