



Gramin Krishi Mausam Sewa
INDIA METEOROLOGICAL DEPARTMENT
ODISHA UNIVERSITY OF AGRICULTURE & TECHNOLOGY
Mahisapat, Dhenkanal- 759 013



Deepanjali Dugal,
Jr. Agrometeorologist

Week No.02

Bulletin No.04
Dt. 12.01.2024
(FRIDAY)

DISTRICT: DHENKANAL (Mid-central Table Land Zone)

Weather condition during last week (06.01.2024 to 12.01.2024)

Rainfall (mm)	0.0
Maximum Temperature (°C)	28.9-30.0
Minimum Temperature (°C)	13.8-19.1

Forecast (Up to 17.01.2024) given by Met. Centre, IMD, Bhubaneswar

Date	Rainfall (mm)	T-MAX (°C)	T-MIN (°C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
13-01-2024	0	27	16	0	73	33	5	135
14-01-2024	0	27	15	0	77	25	6	158
15-01-2024	0	27	15	0	85	26	5	144
16-01-2024	0	28	16	0	92	32	6	150
17-01-2024	0	28	17	4	92	36	9	154

As per IMD forecast, the Dhenkanal district is likely to dry for the next five days with clear to partly cloudy sky. The wind speed is likely to remain 05-09 km/h. The maximum temperature is likely to remain between 27-28°C and minimum temperature between 15-17°C. The maximum RH will remain between 73% to 92% and minimum RH between 25% to 36% for the next five days.

Agromet Advisory

General Advisory:

- ❖ Transplanting of 25-30 days old Paddy seedlings should be done at a spacing of 20 x15 cm, plant 2-3 seedlings per hill for high yielding varieties.
- ❖ In Zinc deficient soil apply Zinc Sulphate (21%) @ 10 kg/acre at every three years.
- ❖ To control bacterial blight in Paddy, dip the seedling roots in 0.1% Plantomycin solution before transplanting for 30 minutes.
- ❖ Drain out excess water from paddy field before herbicide application and irrigate the field after two days of application of herbicide.

PADDY:

Fertilizer Application-

Farmers those who are going for transplanting should apply 35kg DAP, 30kg MOP and 8kg Urea per acre during last puddling. For sandy soil apply 35 kg DAP, 15 kg MOP and 8 kg Urea per acre during last puddling.

Weed management-

In transplanted and broadcasted paddy to control grasses, sedges and broad leaf weeds farmers are advised to spray, Bispyribac Sodium 10% E.C. @ 80 ml/acre or Ethoxysulfuron 15% WDG @ 40g/ acre at 3 to 4 leaf stage by mixing in 200 litre of water.

BLACKGRAM AND GREENGRAM:

Due to foggy weather there is a chance of powdery mildew disease in Black gram and Green gram. To control this disease, spray Sulphur 80% WP @800 g/acre or Hexaconazole 5% E.C @ 300 ml/acre.

MUSTARD:

GROUNDNUT:

White grub infestation in groundnut can be minimized by drenching Chloropyriphous @ 6ml/lit of water or Imidacloprid 30.5 sc@ 1ml/lit of water at the plant base.

COLE CROPS:

Prevailing weather is favorable for attack of tobacco caterpillar in Cole crops. To manage this pest biologically, use Pheromone trap Spodolure @ 6/acre. Instead of this install light trap @ 8 nos. Apply Bacillus thuringensis @ 400 g/acre or Beauveria bassiana @ 500 g/acre. For chemical control, spray Chloropyriphos 50 % + Cypermethrin 5 % E.C @ 200-ml/acre or Lambda Cyhalothrin 5% EC @ 300 ml/acre during afternoon hours by mixing with 200 litres of water.

POTATO:

For controlling Aphids in Potato during primary stage of pest infestation spray Neem Based Pesticide (Azadirachtin) 1500 PPM @ 600-ml/acre. For chemical control spray Thiamethoxam 25% W.G @ 40-gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water.

TOMATO:

To control early and late leaf blight in tomato, spray Mancozeb 75% WP @ 600 g/acre or Zineb 75% WP @ 500gram/acre at 7 days interval.

BRINJAL:

Use yellow sticky traps 5 @ 8-10/acre to attract and kill White fly in Brinjal. During primary stage of pest infestation spray neembased pesticide (Azadirachtin) 1500 PPM @ 600-ml/acre by mixing in 200-litre of water. To control this pest chemically spray Thiamethoxam 25 % WG @ 40 g/acre or Pyriproxifen 10% EC @ 200 ml/acre or Spiromesifen 22.9% SC @ 250 ml/acre by mixing in 200 litre of water.

ONION:

Onion Seedlings of 8-weeks old which are 12-15 cm in height should be planted in the main field. Before transplanting apply 10 tonnes of Compost, 52 kg of DAP, 30 kg of Murate of Potash 20 kg of urea and 20 kg of sulphur and mix it to the soil uniformly. Transplant the seedlings in line by keeping the row to row spacing at 25 cm and plant to plant spacing at 10 cm.

MANGO:

To control mango hopper, spray Imidacloprid @60ml or Carbendazim @200g/200lit of water.

Approved by ADR cum Nodal Officer
GKMS-AMFU, RRTTS
Mahisapat, Dhenkanal

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>),

(iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)