



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Ranital, Bhadrak, - 756111

WeekNo.02

No.-04 (Eng.Bulletin)
Dt.-12.01.2024

District-Jajpur (North-Eastern Coastal Plain Zone)

Forecast (Upto 17.01.2024)
Given by Met. Centre,IMD,Bhubaneswar

DISTRICT: Jajpur– As per IMD forecast, the district is likely to receive **Dry weather for 5 days** with Clear sky to partly cloudy sky. The wind speed is likely to remain within 05-08 km/h. The maximum temperature and minimum temperature will likely to remain 27-28°C and 16-17°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
JAJPUR	13-01-2024	0	27	16	0	53	31	6	112
	14-01-2024	0	27	16	0	68	22	6	202
	15-01-2024	0	28	16	0	65	24	5	236
	16-01-2024	0	28	17	0	81	31	5	112
	17-01-2024	0	27	17	4	86	39	8	158

For further information, contact the Met.Centre,Aerodrom Area,IMD,Bhubaneswar,Tel.#0674-2596116.

Agromet Advisory

Kharif Rice	<ul style="list-style-type: none"> ❖ For storage of Rice make the storage godown clean and spray Lambda Cyahalothrin <u>4.9CS@ 1.5ml/li.</u> of water to minimize the chance of future infestation of storage pest.
Blackgram	<ul style="list-style-type: none"> ❖ After harvesting of paddy, it is ideal time for land preparation for sowing pulses like green gram, black gram and pea. ❖ Sow the seeds after treating with (Carboxin+Thiram)@2gm/kg of seed. ❖ Treat the seed after 7 days of above insecticide with specific Rhizobium culture before sowing. Mix 400g culture with 3g Ammonium molybdate & 800 ml water and apply with 10 kg of seed. ❖ Seed rate: Green gram and Black gram: 25-30Kg/ha. Pea-50Kg/ha. Fertilizer dose:20-40-20 Kg N-P2O5-K2O/ha
Rapeseed and Mustard	<ul style="list-style-type: none"> ❖ Thinning should be carried out before top dressing to maintain optimum plant population.Perform hoeing, weeding and thinning (wherever necessary) at 3 weeks stage. ❖ Irrigate the field if soil moisture stress is observed. ❖ Past weather condition is favourable for aphid attack and Leaf spot incidence in mustard crop so its regular monitoring should be advised. ❖ To manage leaf Webber,Aphid and Sawfly pests at early stage of infestation spray Neem-based pesticide (Neem Oil 1500 PPM) @ 600-ml/acre by mixing in 200-litre of water. After 7-10 days need based spraying of Acephate @300g/acre is recommended.
Vegetables	<p style="background-color: yellow; margin: 0;">Cole Crops :</p> <ul style="list-style-type: none"> ❖ To prevent collar rot of cole crops Spraying of (Metalaxyl+Mancozeb) @400g/acre or (Cymaxonil+Mancozeb) @ 2g /lit of water recommended. ❖ To manage Tobacco Caterpillar,Cabbage webber & Cabbage head borer spray Neem based formulation 1500ppm @600ml/acre followed by neem based spraying of Examectin Benzoate 5SG (80-100)g/acre or spinosad 45SC@70ml/acre recommended. ❖ In Boron deficient soil foliar spray of 20% Boron @300g/acre to prevent the symptoms of hollow stems is recommended. <p style="background-color: yellow; margin: 0;">Tomato & Brinjal:</p> <ul style="list-style-type: none"> ❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base. ❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants. ❖ For minimizing the infestation of sucking insect pests like white fly, leaf hopper,thirps and aphid in

	<p>vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water or Difenthurian @1g/li Of water.</p> <ul style="list-style-type: none"> ❖ To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly & tomato apply 0.5 g Emamectin benzoate or Spinosad 0.4 ml/ liter of water. <p>Cucurbits:</p> <ul style="list-style-type: none"> ❖ Sowing of Recommended HYV/Hybrid of Cucurbits after treating the seeds with Imidachloprid 600FS@(8-10)ml/kg seeds to minimize the incidence of viral /phytoplasma incidence.
Flower	<ul style="list-style-type: none"> ❖ Marigold ❖ Spray the crop with Difenthurion @1g/li. Of water or Abamectin @0.75ml/li. Of water to control the infestation of Thrips,Aphids,Mites and Leaf Minor. ❖ For spraying of incidence of leaf spot & Blight spraying of (Carbendazim+Mancozeb) Recommended.
Fruit	<ul style="list-style-type: none"> ❖ Papaya-Install Yellow Sticky trap to trap the vector (Whitefly) of Papaya Leaf Curl Virus. Alternate spraying of Neem oil formulation 1500ppm @3ml/lit of water and flonicamid 50WG @0.3g/litre to control whitefly. ❖ Coconut- ❖ To reduce the incidence of Leaf blight in newly planted coconut saplings spraying of Meatalxyl+ Mancozeb@2 g/ litre of water is advised.
Pisciculture	<ul style="list-style-type: none"> ❖ As the winter is in the starting phase, there is a possibility of “EUS” disease in fish. Farmers are advised to apply ‘Cifax’ @ 400 ml mixed with 40 litre of water in one acre pond area as a preventive measure. ❖ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal Science	<ul style="list-style-type: none"> ➤ The large ventilation spaces in shed must be closed in night, as the night temperature is falling. ➤ Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals

Mrs. Meera Mahanty
Jr. Agrometeorologist
AMFU, RRTTS, Ranital

Dr. Umasankar Nayak
ADR & NODAL OFFICER
AMFU, RRTTS, Ranital

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides,Vermi &Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download:(Android:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>),(iOS:<https://apps.apple.com/in/app/meghdoot/id1474048155>)