

GRAMIN KRISHI MAUSAM SEWA



INDIA METEOROLOGICAL DEPARTMENT NORTH EASTERN GHAT ZONE

RRTTS, G. UDAYAGIRI, OUAT, KANDHAMAL, ODISHA-762100 MULTIMODEL ENSEMBLE BASED DISTRICT LEVEL WEATHER FORECAST FOR THE STATE: ODISHA

ISSUED ON: 12.01.2024 VALID TILL 08:30 IST OF THE NEXT 5 DAYS

Sri Rahul Kumar Sahu Jr. Agrometeorologist Week No. 02

Dt: 12.01.2024 Bulletin No.04

Forecast (up to 17.01.2024) Given by Met. Center IMD, Bhubaneswar

DISTRICT: KANDHAMAL – As per the IMD forecast, the district is likely to receive no rainfall in next five days with a clear to mainly clear sky. The wind is likely to remain within 5 -6 km/hr. The maximum temperature and minimum temperature will likely to remain between 27-28°C and 11-13°C respectively. The Maximum Relative Humidity and Minimum Relative Humidity will likely to remain between 57-94% and 20-30% respectively for next five days.

DISTRICT	Date	Rainfall (mm)	T- MAX (C)	T- MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
PHULBANI	13-01-2024	0	27	12	0	70	30	5	169
	14-01-2024	0	27	11	0	57	25	6	195
	15-01-2024	0	27	11	0	66	20	5	161
	16-01-2024	0	28	12	0	91	25	5	199
	17-01-2024	0	27	13	4	94	29	6	163

Agromet Advisory

- Pea thrips may be seen sucking the flowers, stems. To control this, apply Roger 0.03% or Malathion 0.1% to the crop.
- For Rice, use varieties such as Lalat, Swarna Samrudhi, Kalinga Dhan, Mandakini, Manaswini.
- Sow pre-germinated treated paddy seeds in puddled soil either by broadcasting or by using drum seeder.
- Maintain only a thin film of water for proper establishment and early growth of Rice seedlings.
- During cool winter days to protect the rice nursery from cold injury.
- To make warm environment use a greater number of light bulbs to protect the chicks in winter.
- Increase the feeding time for chicks and provide clean water
- If chicks are congregated at one place, then provision for more warmth.
- Close the open spaces in night and early morning to protect them winter

GARDEN PEA:

- Stage- flowering in late sown crop
- Pea thrips may be seen sucking the flowers, stems and pods. To control this, apply Roger 0.03% or Malathion 0.1% to the crop.
- You can also use Azadirachtin 0.03% or NSKE 5% in addition.



Pea thrips attack

RICE:

- Stage- nursery
- Use varieties such as Lalat, Swarna Samrudhi, Kalinga Dhan, Mandakini, Manaswini.
- Sow pre-germinated treated paddy seeds in puddled soil either by broadcasting or by using drum seeder.
- Maintain only a thin film of water for proper establishment and early growth of Rice seedlings.
- During cool winter days to protect the rice nursery from cold injury.
- Apply a thin layer of well decomposed FYM after sowing of seeds in wet bed to keep the bed relatively warm.

CABBAGE -

- To control diamond back moth in cabbage, install 6-8 light traps per acre.
- During primary stage of pest infestation spray neem-based pesticide @ 1-litre/acre by mixing in 200-litre of water.
- If infestation prevails, spray Tolfenpyrad 15 % EC @ 400-ml/acre or Emamectin Benzoate 5 % SG @ 80-gram/acre by mixing in 200-litre of water.

TOMATO –

- Stage- early vegetative
- If fruit borer infestation prevails, Spray any one of the insecticides Chlorantraniliprole 18.5% S.C @ 80 ml/acre or Flubendiamide 39.5 % M/M S.C @ 40 ml/acre or Indoxacarb 14.5% S.C. @ 200 ml/acre.
- Use pheromone trap 20@acre.

POULTRY:

- To make warm environment use a greater number of light bulbs to protect the chicks in winter.
- Increase the feeding time for chicks and provide clean water
- If chicks are congregated at one place, then provision for more warmth.
- Close the open spaces in night and early morning to protect them winter.
- If any disease arises, isolate the affected one and vaccinate the others.

For better agriculture benefits use, OUAT KALINGA products (Seedling, Sampling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value added products, Fingerlings/Yearlings, Poultry chicks, Breeds, Mushroom and Spawns, Bio-fertilizers, Bio-pesticides, Vermi & Vermi Compost).

ASSOCIATE DIRECTOR OF RESEARCH & NODAL OFFICER