



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Bhubaneswar -751 003

Dr.T.R Mohanty
Nodal Officer

Week No.02

No. – 35 (Eng. Bulletin)
Dt. -09.01.2024

District – Puri (East and South-Eastern Coastal Plain Agroclimatic Zone)

The mean maximum daily temperature was 27.57°C and mean minimum daily temperature was 17.28°C of the Puri district during the last week. The district received no rainfall during the last seven days. Harvesting of Long duration paddy is going to be completed. Raising of nursery bed and land preparation for rabi paddy is going on. Harvesting and planting of Sugarcane is under progress. Sowing of Rabi crops like Cereals, Pulses, Oilseeds, Vegetables and Spices are under progress. Crop coverage up to the end of this week is 48%. Overall crop condition is Normal.

Forecast (Up to 14.01.2024)

Given by Met. Centre, IMD, Bhubaneswar

DISTRICT: PURI – As per the forecast received from IMD, dry weather may prevail over the district of Puri for next five days. The sky condition may remain clear to mainly clear for the next five days. The wind may flow North Easterly to South Easterly with speed of 09-10 km/h. The daily maximum and minimum temperature are likely to remain 26-28°C and 16-19°C for next five days.

DISTRICT	PURI				
Date	10/01/2024	11/01/2024	12/01/2024	13/01/2024	14/01/2024
Rainfall (mm)	0	0	0	0	0
T-MAX (C)	27	28	27	26	26
T-MIN (C)	18	19	18	17	16
Cloud Cover	2	2	0	0	0
Rh Max (%)	78	72	77	79	77
Rh Min (%)	47	52	54	51	45
Wind speed (kmph)	10	9	10	10	9
Wind Direction (deg)	112	153	120	109	161

For further information, contact the Met. Centre, Aerodrom Area, IMD, Bhubaneswar, Tel. # 0674-2596116.

Agromet Advisory

- ❖ Maintain saturated condition in the surface soil of nursery bed by applying irrigation in furrow, for proper establishment and early growth of Rice seedlings.
- ❖ Mulch vegetable seedling with straw or other mulching material to avoid cold injury.
- ❖ In view of prevailing low temperature, Provide light irrigation to standing vegetables and crops during morning hours.
- ❖ Farmers are advised to prepare nursery bed and treat the paddy seeds with Carbendazim 50% W.P @ 1.5 g/1kg of seeds or Carboxin 37.5% + Thiram 37.5 % @ 3 g /1 kg of seeds before sowing in the nursery.
- ❖ Standing water to a depth of 2-3 cm should be maintained at least for 2-3 days before uprooting of seedlings.
- ❖ Sunflower crop is at 25-30 days stage. Go for hoeing and weeding and thinning and apply N -fertilizer.
- ❖ Spraying of pesticides to be done after 3 pm to protect the honey bee.
- ❖ Cover the recently planted cole crops with paper pockets during night time to prevent from low temperature stress.
- ❖ For maintaining the heat in the poultry shed during winter season, install heaters in the shed.

PADDY (Summer): Nursery Management

Prepare the wet nursery seed beds for paddy. Irrigate the nursery field and puddle it 2-3 times followed by planking. Apply 200 kg of FYM, 4 kg of DAP, 2.5 kg of MOP and 1 kg of Zinc sulphate during last puddling. Divide nursery area into smaller plots of convenient length. Irrigation channels of size 30 cm in width should be made along the beds for irrigation/drainage purpose. Sprouted Seeds should be sown @ 40-50 g / m² area of seed bed by line sowing with 5 cm gap between each line.

Seed treatment

Treat the paddy seeds with Carbendazim 50% W.P @ 1.5 g/1kg of seeds or Carboxin 37.5% + Thiram 37.5 % @ 3 g /1 kg of seeds before sowing in the nursery.

SUGARCANE: To prevent red rot disease, resistant varieties like CO-7219, CO-8371, CO-8013, CO-8014, CO-6907, CO-7601 and CO- 7602 should be selected. Soak the setts for 30 minutes in a solution of 100 lit of water containing 150 g of carbendazim 50 WP, 200- ml of chlorpyrifos 20 % EC and 1 kg urea. Do not add chlorpyrifos if soil drenching is adopted.

CHILLI:-Mite

For controlling mites in chilli, spray Ethion 50 % EC @ 600 ml/acre or Spiromesifen 22.9 % SC @ 200 ml/acre or Propargite 57 % EC @ 600 ml/acre.

POTATO: Go for hoeing and hand weeding in Potato after 3 weeks of first earthing up. Apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 400 ml/acre for control of cut worm and leaf eating insect.

CABBAGE: Cabbage webber and tobacco caterpillar -

Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in Cabbage. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water.

ROSE: To control thrips in rose, spray Fenpyroximate 5% EC @ 2 ml or Abamectin @1 ml per litre of water.

ANIMAL HUSBANDRY: If the animals have not yet been vaccinated against FMD, Haemorrhagic Septicaemia, Black Quarter, Enterotoxemia etc., these must be done. Due care should be taken to store/procure fodder for periods of shortage that may occur during the winter months in certain areas. Perennial grasses must be cut at this time.

PRINCIPAL NODAL OFFICER

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>) (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)