

Gramin Krishi Mausam Sewa India Meteorological Department Orissa University of Agriculture and Technology, Regional Research & Technology Transfer Sub Station,



Kirei, Sundergarh-770073

Dr. Sajeeb Kumar Biswasi, Senior Scientist & Nodal Officer, GKMS No-03(Eng.Bulletin) *Dt* -09.01.2024

Week No - 02

District –Sundergarh (North Western Plateau Agroclimatic Zone)

The mean maximum daily temperature was 27.2°C & mean minimum daily temperature was 15.6°C in Sundergarh district during last week. The District received in average 0.0mm rainfall during the last week. The kharif Paddy is at maturity to harvesting stage. Early sown non-Paddy crops like Pulses, Oilseeds crop are at harvesting stage. The Vegetables are at harvesting stage. The overall crop condition is normal.

<u>Forecast (Up to 14.01.2024)</u> Given by Met.Centre, IMD, Bhubaneswar

DISTRICT: SUNDERGARH- As per IMD Forecast, the district is likely receive no rainfall for the next five days. The sky condition is clear sky to mainly clear for the next five days. The wind speed is likely to remain within 5-7 km/h. The Maximum Temperature and Minimum Temperature is forecasted to remain between 26-28°C and 11-15°C respectively for the next five days.

DISTRICT					
Date	10/01/2024	11/01/2024	12/01/2024	13/01/2024	14/01/2024
Rainfall (mm)	0	0	0	0	0
T.MAX(°C)	26	27	28	27	27
T.MIN (°C)	14	15	13	12	11
Cloud Cover	2	2	0	0	0
Rh Max (%)	64	64	56	36	39
Rh Min (%)	31	31	35	20	15
Wind speed	5	6	7	6	6
(kmph)					
Wind Direction	71	72	71	101	109
(deg)					

For further information, contact the Met. Centre, Aerodrom Area, IMD, Bhubaneswar, Tel.# 0674-2596116

Agromet Advisory

DISTRICT:SUNDERGARH:-

Paira Crop: The pulses grown as of paira crop to be sprayed with 2% DAP solution for crop vigour.

Rabi Paddy: Chose short and medium duration paddy variety for rabi season. Complete the nursery preparation in 10 decimel area for one acre. For this the beds are to be prepared 3 ft width with suitable length and leave a space in between the beds for facilitating the intercultivation. In the beds apply add 40 baskets of cow dung, 12 kg of single super phosphate and 2 kg of murate of potash.

Soak the seeds in water for 24hours and after germination, sow them in the beds.

Sweetpotato:

At 25 to 30 days old, weed the crop and fertilize it with 37.5 kg of potassium per acre and 37.5 kg of potash fertilizer.

Irrigate by looking at the soil moisture.

Wheat

Irrigate the crop at crown root initiation and jointing stage. In late sowing go for intercultivation. In the early sown crop apply 8 kg Nitrogen per acre.

Groundhut Prepare the land for groundnut cultivation. Chose the sandy loam soil having irrigation facility. The winter groundnut at 35-40 days after sowing do not disturb the soil due to peg formation stage.

Sunflower:

At 25 to 30 days after sowing the sunflower, apply 6-8 kg nitrogen per acre. Go for thinning operation.

Toria:

Due to powdery mildew, apply 1 kg sulphur powder in 200 liters of water per acre.

Potato: In the delayed sown crop, go for intercultivation and earthing up and apply the remaining fertilizer of Nitrogen 30 kg/acre.

Animal: The shed should be well ventilated.

It is best to put sand on the floor of the shed because their excreta is absorbed into the sand so they can always stay on a dry floor.

Always keep inside and outside of the cowshed clean

In winter, cattle are fed with high-energy fiber foods such as grass straw, straw, sorghum stalks and , corn leaves.

Poultry:

For more eggs and meat, keep Banraja, Grampriya breeds of chickens.

Vaccinate chickens regularly against distemper and vernal distemper as advised by a veterinarian.

For any problem related to agriculture sector, please dial the toll free number 1551.

For better agricultural benefits use, **OUAT KALINGA** Products (Seedlings, Saapplings and QPM, Tissue culture plants, Fruits and vegetables, value added products, Fingerlings/Yearlings, Poultry chicks, Breeds, Mushrooma and spwn, Biofertilizer, Biopesticides, Vermi & Vermicompost and Farm implements)

IMD weather forecast & Agrometeorological Advisory of Odisha location is now available on Meghdoot mobile app in English & Odia Language.

Download: (Andriod: https://play.google.com/store/apps/details?id=com.aas.meghdoot)

ବି : ଦୃଷ୍ଟବ୍ୟ :ଓୟୁଏଟି ଶାଖା ଚିପଲିମା ଓ ଗୟାରିପାଲି ବିହନ ଗବେଷଣା ଫାର୍ମ ଠାରେ ସ୍ପର୍ଶ ଧାନ ବିହନ ୨୦ କିଗ୍ରା ବଞା ପିଛା ଟ. ୭୧୨ ୪୦ . ପଇସା ଚାଷୀ ନିମିତତ ବିକରି ହଉେଅଛି ।

Nodal Officer