



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Ranital, Bhadrak, - 756111

WeekNo.05

No-10(Eng.Bulletin)
Dt.-02.02.2024

District-Balasore (North-Eastern Coastal Plain Zone)

Weather Condition During Last Week at Balasore (27.01.2024 –02.02.2024)

Rainfall (mm)	0.5
Maximum temperature (⁰ C)	25.0-28.0
Minimum temperature (⁰ C)	13.0-20.0

Forecast (Upto 07.02.2024)

Given by Met.Centre,IMD,Bhubaneswar

DISTRICT: Balasore– As per IMD forecast, the district is likely to receive **Dry weather for 3 days & Very light to light rain for 2 days** with clear sky to Overcast Sky. The wind speed is likely to remain within 09-15km/h. The maximum temperature and minimum temperature will likely to remain 26-29°C and 19°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
BALASORE	03-02-2024	4	26	19	8	78	51	11	180
	04-02-2024	0	28	19	2	82	41	10	150
	05-02-2024	0	29	19	0	83	31	9	117
	06-02-2024	1	29	19	0	87	41	10	199
	07-02-2024	0	28	19	0	90	56	15	199

For further information,contact the Met.Centre,Aerodrom Area,IMD,Bhubaneswar,Tel.#0674-2596116.

Agromet Advisory

Rice	<ul style="list-style-type: none"> ❖ Utilising the available water (Rainfall/Irrigation) puddle the land &complete the transplanting of summer rice with basal application of 35kg DAP,20 kg MOP &10 kg ZnSO₄/acre. ❖ After the unseasonal rainfall there is increase in humidity in the storage which may lead to infestation of insect pest & occurrence of disease. ❖ Inspect the go down &open the doors & windows for sometime for ventilation & need based spraying of Lambda Cyahalothrin<u>4.9CS@1.5ml/li.</u> of water to minimize the infestation of storage pest.
Blackgram	<ul style="list-style-type: none"> ❖ To reduce incidence of Powdery mildew, Anthracnose & Stem rotinblackgram spraying of (Tebuconazole + Sulphur) @500g/acre recommended.
Rapeseed & Mustard	<ul style="list-style-type: none"> ❖The mustard crop is at the flowering stage. The rainfall received during the last week which results in higher humidity and lower temperature might aggravate Aphid, Leaf Webber and Tobacco Caterpillar infestation in mustard crop. To manage these pests at early stage of infestation spray Neem-based pesticide (Neem Oil 1500 PPM) @ 600-ml/acre by mixing in 200-litre of water. After 7-10 days need based spraying of Emamectin Benzoate @300g/acre is recommended. ❖Spraying of Insecticide should be done at the afternoon hours to protect the honeybees & other beneficial insects.
Greengram	<ul style="list-style-type: none"> ❖Complete the sowing of summer Greengram as soon as possible to avoid the infestation of sucking pest and viral disease under delayed sowing condition.
Vegetables	<p>Tomato & Brinjal:</p> <ul style="list-style-type: none"> ❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base. ❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants. ❖ For minimizing the infestation of sucking insect pests likewhite fly, leaf hopper,thirps and aphid in vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water or

	<p>Difenthurian @ 1g/li Of water.</p> <ul style="list-style-type: none"> ❖ To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly & tomato apply 0.5 g Emamectin benzoate or Spinosad 0.4 ml/ liter of water. <p>Cucurbits:</p> <ul style="list-style-type: none"> ❖ Sowing of Recommended HYV/Hybrid of Cucurbits after treating the seeds with Imidachloprid 600FS@(8-10)ml/kg seeds to minimize the incidence of viral /phytoplasma incidence.
Flower	<ul style="list-style-type: none"> ❖ Marigold ❖ Spray the Marigold plant with Difenthurion @ 1g/l Of water or Abamectin @0.75ml/l Of water to control the infestation of Thrips,Aphids,Mites and Leaf Minor. ❖ For spraying of incidence of leaf spot & Blight spraying of (Carbendazim+Mancozeb) recommended.
Fruit	<ul style="list-style-type: none"> ❖ Papaya-Install Yellow Sticky trap to trap the vector (Whitefly) of Papaya Leaf Curl Virus.Alternate spraying of Neem oil formulation 1500ppm @3ml/l of water and flonicamid 50WG @0.3g/l to control whitefly.
Pisciculture	<ul style="list-style-type: none"> ❖ As the winter is in the starting phase, there is a possibility of “EUS” disease in fish. Farmers are advised to apply ‘Cifax’ @ 400 ml mixed with 40 litre of water in one acre pond area as a preventive measure. ❖ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal Science	<ul style="list-style-type: none"> ➤ The large ventilation spaces in shed must be closed in night, as the night temperature is falling. ➤ Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals .

Mrs. Meera Mahanty
Jr. Agrometeorologist
AMFU, RRTTS, Ranital

Dr. Umasankar Nayak
ADR &NODAL OFFICER
AMFU,RRTTS, Ranital

For better agricultural benefits use, **OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides,Vermi&VermiCompostandFarmImplements).**

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download:(Android:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>),(iOS:<https://apps.apple.com/in/app/meghdoot/id1474048155>)