

Gramin Krishi Mausam Sewa India Meteorological Department Odisha University of Agriculture and Technology Ranital, Bhadrak, - 756111

No-10(Eng. Bulletin) Dt.-02.02.2024

District-Bhadrak (North-Eastern Coastal Plain Zone)

WeekNo.05

Weather Condition During Last Week at Bhadrak (27.01.2024 –02.02.2024)

Rainfall (mm)	0.0
Maximum temperature (⁰ C)	26.0-30.0
Minimum temperature (⁰ C)	14.0-19.0

Forecast (Upto 07.02.2024) Given by Met.Centre,IMD,Bhubaneswar

DISTRICT: Bhadrak—As per IMD forecast, the district is likely to receive **Dry weather for 2 days &Very light rain for 3 days** with clear sky to Overcast Sky. The wind speed is likely to remain within 08-15km/h. The maximum temperature and minimum temperature will likely to remain 28-31°C and 19°C to respectively for the next five days.

immum temperature will likely to remain 2				11 20 31 0	20 91 C und 19 C to respectively for the next five days.						
	DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)	
	BHADRAK	03-02-2024	2	28	19	8	85	38	11	198	
		04-02-2024	0	30	19	0	93	45	9	292	
		05-02-2024	1	31	19	0	89	41	8	135	
		06-02-2024	0	30	19	1	88	47	10	112	
		07-02-2024	2	30	19	0	90	46	15	198	

For further information, contact the Met.Centre, Aerodrom Area, IMD, Bhubaneswar, Tel. #0674-2596116.

Agromet Advisory

	Agromet Advisory				
	❖ Utilising the available water (Rainfall/Irrigation) puddle the land &complete the transplanting				
	of summer rice with basal application of 35kg DAP,20 kg MOP &10 kg ZnSO ₄ /acre.				
Rice	❖ After the unseasonal rainfall there is increase in humidity in the storage which may lead to				
	infestation of insect pest & occurrence of disease.				
	❖ Inspect the go down & open the doors & windows for sometime for ventilation & need based				
	spraying of Lambda Cyahalothrin <u>4.9CS@1.5ml/li</u> . of water to minimize the infestation of				
	storage pest.				
Blackgram	❖ To reduce incidence of Powdery mildew, Anthracnose & Stem rotinblackgram spraying of				
	(Tebuconazole + Sulphur) @500g/acre recommended.				
Rapeseed & Mustard	❖The mustard crop is at the flowering stage. The rainfall received during the last week which				
	results in higher humidity and lower temperature might aggravate Aphid, Leaf Webber and				
	Tobacco Caterpillar infestation in mustard crop. To manage these pests at early stage of				
	infestation spray Neem-based pesticide (Neem Oil 1500 PPM) @ 600-ml/acre by mixing in				
	200-litre of water. After 7-10 days need based spraying of Emamectin Benzoate @300g/acre is				
	recommended.				
	❖Spraying of Insecticide should be done at the afternoon hours to protect the honeybees & other				
	beneficial insects.				
Greengram	Complete the sowing of summer Greengram as soon as possible to avoid the infestation of				
	sucking pest and viral disease under delayed sowing condition.				
	Tomato & Brinjal:				
	❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake				
Vegetables	and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base.				
, ogetables	❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants.				

	 For minimizing the infestation of sucking insect pests likewhite fly, leaf hopper,thirps and aphid in vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water or Difenthurian @1g/li Of water. To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly & tomato apply 0.5 g Emamectin benzoate or Spinosad 0.4 ml/ liter of water. Cucurbits:
	Sowing of Recommended HYV/Hybrid of Cucurbits after treating the seeds with Imidachloprid 600FS@(8-10)ml/kg seeds to minimize the incidence of viral /phytoplasma incidence.
Flower	❖ Marigold
	❖ Spray the Marigold plant with Difenthurion @1g/l Of water or Abamectin @0.75ml/l Of water to control the infestation of Thrips, Aphids, Mites and Leaf Minor.
	❖ For spraying of incidence of leaf spot & Blight spraying of (Carbendazim+Mancozeb) recommended.
Fruit	Papaya- Install Yellow Sticky trap to trap the vector (Whitefly) of Papaya Leaf Curl Virus. Alternate spraying of Neem oil formulation 1500ppm @3ml/l of water and flonicamid 50WG @0.3g/l to control whitefly.
Pisciculture	❖ As the winter is in the starting phase, there is a possibility of "EUS" disease in fish. Farmers are advised to apply 'Cifax' @ 400 ml mixed with 40 litre of water in one acre pond area as a preventive measure.
	❖ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal Science	➤ The large ventilation spaces in shed must be closed in night, as the night temperature is falling.
	Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals .

Mrs. Meera Mahanty Jr. Agrometeorologist AMFU, RRTTS, Ranital Dr. Umasankar Nayak ADR &NODAL OFFICER AMFU,RRTTS, Ranital

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compostand Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

 $\label{lem:com} \textbf{Download:} (\textbf{Android:} \underline{\textbf{https://play.google.com/store/apps/details?} id = com.aas.meghdoot),} (iOS:\underline{\textbf{https://apps.apple.com/in/app/meghdoot/id1474048155}})$