



**Gramin Krishi Mausam Sewa**  
**India Meteorological Department**  
**Odisha University of Agriculture and Technology**  
**Ranital, Bhadrak, - 756111**



WeekNo.09

No-17(Eng.Bulletin)  
Dt.-27.02.2024

District-Bhadrak (North-Eastern Coastal Plain Zone)

**Weather Condition During Last Week at Bhadrak (21.02.2024 –27.02.2024)**

Rainfall (mm)	0.0
Maximum temperature (°C)	27.0-36.0
Minimum temperature (°C)	19.0-23.0

Forecast (Upto 03.03.2024)  
Given by Met.Centre,IMD,Bhubaneswar

**DISTRICT: Bhadrak**– As per IMD forecast, the district is likely to receive **Light rain for one day & Dry weather for 4 days** with clear sky to Partly cloudy sky. The wind speed is likely to remain within 10-16 km/h. The maximum temperature and minimum temperature will likely to remain 33-34°C and 20-21°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
BHADRAK	28-02-2024	0	33	20	0	66	35	11	243
	29-02-2024	0	34	21	1	78	30	10	202
	01-03-2024	4	34	21	0	79	39	11	243
	02-03-2024	0	34	21	1	76	31	10	236
	03-03-2024	0	34	21	4	72	27	16	206

For further information, contact the Met.Centre,Aerodrom Area,IMD,Bhubaneswar,Tel.#0674-2596116.

**Agromet Advisory**

<b>Rice</b>	<p><b>General Advisory-</b> As per IMD forecast dry weather is predicted so farmers are advised to-</p> <ul style="list-style-type: none"> <li>❖ <b>Monitor the incidence of various disease pests in field.</b></li> <li>❖ <b>Transplanting of vegetables should be done in afternoon hour to avoid high temperature.</b></li> <li>❖ Transplanting should be advised in case of 21 to 25 days or above old seedlings. For weed management in rice Application of granular herbicide Bensulfuron-methyl @ 4 kg/ acre + Pretilachlor at 5-8 days after transplanting or spraying of Pyrazosulfuron-ethyl 10 WP @ 5 g/10 ltrs of water at 3-5 DAT or Bispyribac sodium @ 120 ml by mixing with 140 liters of water per acre at 10-15 DAT should be advised under clear weather condition.</li> <li>❖ In-case of Leaf blast infestation, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 80 g/acre in 200 litres of water. If infestation of Thrips is noticed in rice, spray Azadirachtin 0.15% neem seed kernel based EC formulation @ 800 ml/acre or Lambda-cyhalothrin 5 % EC @ 200 ml/acre.</li> </ul>
<b>Sugarcane</b>	<ul style="list-style-type: none"> <li>❖ In Ratoon crop of sugarcane remove the disease &amp; pest infected plants and dispose the grassy shoot infected plant followed by recommended fertilizer application &amp; spraying of Fipronil 5 Sc @2ml/l of water recommended.</li> <li>❖ Collect setts from the whole cane of a 6–8 month old plant crop free from diseases and insect pests. Select the upper 1/3rd of cane if seeds are collected from 8-12-month-old crop for better sprouting. Sugarcane Sett treatment is a necessary procedure before planting.</li> <li>❖ Sett Rot management -To manage sett rot disease after planting basal drenching with Carbendazim 12% + Mancozeb 63% @ 400-gram/acre or foliar spray with Metalaxyl 8% + Mancozeb 64% WP @ 400-gram/acre by mixing in 200-litre of water.</li> </ul>
<b>Blackgram/ Greengram</b>	<ul style="list-style-type: none"> <li>❖ To control aphids during primary stage of pest infestation spray Neem Based Pesticide (Azadirachtin) 1500 PPM @ 600 ml/acre. Then spray Thiamethoxam 25% W.G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 17.8% SL @ 60-ml/acre or Imidacloprid by mixing in 200 litres of water.</li> </ul>
<b>Rapeseed &amp; Mustard</b>	<ul style="list-style-type: none"> <li>❖ The mustard/Toria crop is at pod filling stage. Aphids in Mustard and Toria can be controlled by spraying Dimethoate 30% EC @ 2 ml/lit. Spraying of Insecticide should be done at the afternoon hours to protect the honeybees &amp; other beneficial insects.</li> <li>❖ Farmers are advised to keep their harvested produce in safe place .</li> </ul>

<b>Vegetables</b>	<p><b>Tomato &amp; Brinjal:</b></p> <ul style="list-style-type: none"> <li>❖ Farmers should start sowing of tomato seed for summer cultivation in nursery bed because prevailing temperature suitable for germination of seed.</li> <li>❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base.</li> <li>❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants.</li> </ul> <p><b>Cucurbits:</b></p> <ul style="list-style-type: none"> <li>❖ Spraying Of Acephate @250g/acre along with neem oil formulation 600ml/acre to minimize damage caused by Red Pumpkin Beetle in cucurbits.</li> <li>❖ To manage Epilachna Beetle in cucumber, spray neem-based pesticide (Neem Oil 1500 PPM) Azadiractin @ 600 ml/acre of water at early stage of infestation. To manage Epilacna Beetle chemically spray Chlorpyriphos 20 % EC @ 400ml/acre or Profenophos 50 % EC@400ml/acre by mixing with 200 litre of water.</li> </ul>
<b>Fruit</b>	<p><b>Watermelon</b>-Present weather condition favourable for watermelon cultivation. Use improved varieties like Sugar baby, Arka manik, Lalsona and hybrid varieties like Madhuri, Maya, Sweet Ruby, Black magic and Sugar pack. For 1 acre planting 400-800 gram of seed is required. Treat the seed with carbendazim @ 1g / kg of seed.</p> <p><b>Mango</b>-Prevailing weather conditions are congenial for Mealy bug incidence in mango. To control mealy bug, Spray Dimethoate 30 EC @ 2 ml/liter of water during February- March as per requirements.</p>
<b>Pisciculture</b>	<ul style="list-style-type: none"> <li>❖ Small white rash appear on the body of fish and feels rough to hand are the symptoms of white spot disease in fish. Excessive mucus secret from fish body. To treat this disease in fish bath affected fish for 5 minutes in the 200 litre water which is mix with 500gram of salt, then release to the pond. This procedure continues for 7-10 days.</li> </ul>
<b>Livestock</b>	<ul style="list-style-type: none"> <li>❖ As temperature is increasing day by day hence, Farmers should provide sufficient clean drinking water to animals and make proper shelter and feed management.</li> </ul>

Mrs. Meera Mahanty  
Jr. Agrometeorologist  
AMFU, RRTTS, Ranital

Dr. UmasankarNayak  
ADR &NODAL OFFICER  
AMFU,RRTTS, Ranital

**For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi&Vermi Compost and Farm Implements).**

Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for IMD Weather forecast & Agromet advisory and “Damini” App for Lightning warning. For better agricultural benefits use.

**Download:(Android:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>),(iOS:<https://apps.apple.com/in/app/meghdoot/id1474048155>)**