



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Ranital, Bhadrak, - 756111

WeekNo.06

No-12(Eng.Bulletin)
Dt.-09.02.2024

District-Jajpur (North-Eastern Coastal Plain Zone)

Forecast (Upto 14.02.2024)
Given by Met. Centre,IMD, Bhubaneswar

DISTRICT: Jajpur– As per IMD forecast, the district is likely to receive **Dry weather for 4 days & light rain on 13th** with Clear Sky to Generally cloudy sky. The wind speed is likely to remain within 07-08 km/h. The maximum temperature and minimum temperature will likely to remain 28-30°C and 19-21°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
JAJPUR	10-02-2024	0	29	19	0	47	21	8	90
	11-02-2024	0	30	19	0	51	18	7	72
	12-02-2024	0	29	20	7	50	17	7	90
	13-02-2024	5	28	21	6	58	19	7	248
	14-02-2024	0	30	21	4	64	19	8	161

For further information ,contact the Met. Centre, Aerodrom Area,IMD,Bhubaneswar,Tel.#0674-2596116.

Agromet Advisory

Rice	<p>Farmers are advised to give necessary irrigation to field & vegetable crops.</p> <ul style="list-style-type: none"> ❖ Farmers those have already completed transplanting, apply granular herbicide Bensulfuron-methyl + Pretilachlor @ 4kg/ acre at 5-8 days after transplanting. ❖ Utilising the available water (Rainfall/Irrigation) puddle the land & complete the transplanting of summer rice with basal application of 35 kg DAP,20 kg MOP &10 kg ZnSO₄/acre. ❖ In bacterial blight endemic areas, dip the seedling roots in 0.1% (1g in one litre of water) Plantomycin solution before transplanting for 30 minutes.
Sugarcane	<ul style="list-style-type: none"> ❖ In Ratoon crop of sugarcane remove the disease & pest infected plants and dispose the grassy shoot infected plant followed by recommended fertilizer application & spraying of Fipronil 5 Sc @2ml/l of water recommended. ❖ Sugarcane Sett treatment with (Metalaxyl + Mancozeb)@2g/l Of water is recommended to decrease the incidence of diseases like Red Rot & Wilt.
Blackgram	<ul style="list-style-type: none"> ❖ To reduce incidence of Powdery mildew, Anthracnose & Stem rot in blackgram spraying of (Tebuconazole + Sulphur) @500g/acre recommended. ❖ To control aphids during primary stage of pest infestation spray Neem Based Pesticide (Azadirachtin) 1500 PPM @ 600 ml/acre. Then spray Thiamethoxam 25% W.G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 17.8% SL @ 60-ml/acre or Imidacloprid by mixing in 200 litres of water.
Rapeseed & Mustard	<ul style="list-style-type: none"> ❖ The mustard/Toria crop is at the flowering to pod filling stage. Aphids in Mustard and Toria can be controlled by spraying Dimethoate 30% EC @ 2 ml/lit. ❖ To control Mustard Sawfly, Leaf Webber and Tobacco Caterpillar infestation in mustard crop at early stage spray Neem-based pesticide (Neem Oil 1500 PPM) @ 600-ml/acre by mixing in 200-litre of water. Spraying of Insecticide should be done at the afternoon hours to protect the honeybees & other beneficial insects.
Vegetables	<p>Cucurbits:</p> <ul style="list-style-type: none"> ❖ Sowing of Recommended HYV/Hybrid of Cucurbits after treating the seeds with Imidachloprid 600FS@(8-10)ml/kg seeds to minimize the incidence of viral /phytoplasma incidence. ❖ Spraying Of Acephate @250g/acre along with neem oil formulation 600ml/acre to minimize

	<p>damage caused by Red Pumpkin Beetle in cucurbits.</p> <p>Tomato & Brinjal:</p> <ul style="list-style-type: none"> ❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base. ❖ Provide bamboo or wooden stake for better fruit set and growth of Tomato plants. ❖ For minimizing the infestation of sucking insect pests like white fly, leaf hopper, thrips and aphid in vegetables nursery and main field, spray Thiamethoxam @ 0.4g/litre of water or Difenthurian @ 1g/li Of water. ❖ To control leaf eating caterpillar and fruit borer in winter vegetables like Brinjal, Cole crops, Chilly & tomato apply 0.5 g Emamectin benzoate or Spinosad 0.4 ml/ liter of water.
Pisciculture	<ul style="list-style-type: none"> ❖ The fishes should be fed daily with oil cake bran mixture at the rate of 2-3 % of their total biomass.
Animal Science	<ul style="list-style-type: none"> ➤ Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals .

Mrs. Meera Mahanty
 Jr. Agrometeorologist
 AMFU, RRTTS, Ranital

Dr. Umasankar Nayak
 ADR & NODAL OFFICER
 AMFU, RRTTS, Ranital

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for IMD Weather forecast & Agromet advisory and “Damini” App for Lightning warning. For better agricultural benefits use.

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)