

# Agromet Advisory Services REGIONAL RESEARCH AND TECHNOLOGY TRANSFER STATION, KENDUJHAR

Week no -05

Date- 02.02.2024

## Bulletin for Kendujhar district of North Central Plateau Agroclimatic Zone.

The mean weekly maximum and minimum temperature at Kendujhar was 21.8 <sup>o</sup>C and 12.2 <sup>o</sup>C respectively with 27.2 rainfall during the last week.

# Weather Forecast (up to 07.02.2024)

There is likely to be **very light rainfall** during the next five days. The wind speed may remain within 5-8 kmph. The maximum and minimum temperature will remain within 27-28  $^{\circ}$ C and 15-17  $^{\circ}$ C during the next five days.

	Date	Rainfall (mm)	T- MAX (C)	T- MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
KENDUJHARGARH	03-02-2024	1	27	17	4	66	41	6	68
	04-02-2024	0	27	16	0	81	29	5	124
	05-02-2024	0	28	15	0	91	26	6	112
	06-02-2024	0	27	16	2	95	41	8	146
	07-02-2024	0	27	17	4	91	37	8	198

**General advisory:** According to IMD, **very light rainfall** over the Keonjhar district is forecasted.

- Harvest the matured crops and store them in safe storage.
- Use Aluminium Phosphide (Celphos tablets) as fumigants for safe storage.
- Carefully, monitor the field for disease and pests.

# **Chickpea:**

- Alternaria blight in chickpea can be managed by spraying Mancozeb @ 2.5 g/lt of water or Carbendazim @ 1 g/lt of water.
- Avoid excessive irrigation to the crop.
- To control powdery mildew disease in chickpea spray with Dithane- M 45 @ 2.5 g/lt of water.

# Linseed:

- In case of Utera/Paira cropping, apply top dressing of Urea in the pre-flowering stage.
- Spraying of Imidachloprid @ 2 g/lt of water can reduce the aphids population.
- Use Chloropyriphos and neem extract containing Azadiractin 300 ppm can reduce the damage by Termite and leaf miner.

#### **Mustard:**

- Mustard are in flowering and pod filling stage.
- Aphids in Mustard and Toria can be controlled by spraying Dimethoate 30% EC @ 2 ml/lit.

### Vegetables:

## • Brinjal:

- Whitefly and aphids in brinjal can be managed by spraying Phosphamidon 40% SL @ 1.5 ml/lt of water.
- For fruit and shoot borer management in brinjal, spray Emamectin benzoate @ 4 g/10lt of water.
- Little leaf in brinjal can be controlled by spraying Methyl demeton 30 EC @ 1 ml/lt of water to control the vectors.

## • Cauliflower:

- Diamond back moth in cauliflower can be managed by spraying Cartap hydrochloride @ 1 g/lt of water.
- Fall army worm or head borer can be controlled by spraying Emamectin benzoate 5% SG @ 4-5 g/10lt of water.
- Aphids in Cauliflower can be controlled by spraying Thiomethoxam 25% W.G @ 80g or Acetamiprid 20% S.P @ 30g per acre in 200 lt of water.

#### • Cabbage:

• To control Cabbage head borer, apply Novaluron 10% EC @ 400 ml or Spinosad 2.5% SC @ 300ml or Emamectin benzoate 5% SG @ 100g in 200lt of water and spray on leaves.

# • Chilli:

• Symptoms of chilli thrips can be seen by leaf crinkles and upward curling. Spray Imidachloprid 17.8% SL @ 3 ml/10 lt of water or Emamectin benzoate 5% SG @ 4g/10 lt of water.

#### • **Tomato:**

- Remove the disease affected leaves and twigs.
- Make the field clean and remove the weeds.
- For managing fruit borer in tomato, spray Flubendiamide 5g/10 lt of water.

#### Flowers:

#### • Rose:

• To control Thrips, aphids and leaf hoppers spray Methyl demeton 25 EC @ 2 ml/lt or Carbofuran 3G @ 5 g/plant.

- Marigold:
- Mealy bug in marigold can be controlled by spraying Fish oil or resin soap @ 25 g/lt.
- In severe infestation, spray Dimethoate or Prophenophos @ 2 ml/lt.

#### Animal Husbandry:

- Cow:
- Provide clean drinking water to the calves and cows.
- Arrange proper heat to the sheds to reduce the cold waves.

#### **TECHNICAL OFFICER**

For better agricultural benefits, use OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.

Download: (Android: https://play.google.com/store/apps/details?id=com.aas.meghdoot), (iOS: https://apps.apple.com/in/app/meghdoot/id1474048155)