



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Ranital, Bhadrak, - 756111



WeekNo.11

No-22(Eng.Bulletin)
Dt.-15.03.2024

District-Balasore (North-Eastern Coastal Plain Zone)

Weather Condition During Last Week at Balasore (09.03.2024-15.03.2024)

Rainfall (mm)	19.6
Maximum temperature (⁰ C)	33.0-36.0
Minimum temperature (⁰ C)	20.0-24.0

Forecast (Upto 20.03.2024)

Given by Met.Centre,IMD,Bhubaneswar

DISTRICT: Balasore- As per IMD forecast, the district is likely to receive **Very Light to Moderate rain for upcoming 5 days** with Partly cloudy Sky to overcast sky.The wind speed is likely to remain within 10-14km/h. The maximum temperature and minimum temperature will likely to remain 29-33°C and 20-25°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
BALASORE	16-03-2024	4	33	20	5	89	52	14	180
	17-03-2024	2	33	20	4	83	49	12	197
	18-03-2024	5	32	24	6	72	50	12	234
	19-03-2024	15	31	25	6	71	41	10	162
	20-03-2024	20	29	25	8	72	40	12	158

For further information,contact the Met.Centre,AerodromArea,IMD,Bhubaneswar,Tel.#0674-2596116.

Crop and Livestock Specific Agromet Advisory

General Advisory	<p>As per IMD, Very Light to Moderate rainfall is expected during next five days ,</p> <ul style="list-style-type: none"> ➤ Farmers are advised to drain excess water from all crops like- Green Gram, Black Gram, Maize, Vegetables & sunflower. ➤ Withhold all spraying & fertilizer application operations. ➤ The runoff water may be properly drained to farm pond for supplemental irrigation during later days. 	
Crop Name	Crop stage	Agromet Advisory
Summer Rice	Transplanting	<ul style="list-style-type: none"> ❖ In summer rice high infestation of stem borer is observed so farmers are advised to monitor the activity of stem borer by installing light trap /Pheromone trap and apply 4 kg Rynaxypyr 0.4 G or 8 kg Fipronil 0.3G per acre.Spraying of Quinalphos @400 ml per acre is advised to prevent egg laying of adults. ❖ In some places in rice field BLB infestation noticed so to control this disease apply 200g plantomycin along with 500 g cupper oxy chloride by mixing 200 l of Water & apply 10 kg MOP /acre.
Sunflower	Flowering	<ul style="list-style-type: none"> ❖ Remove the collar rot infected plants from field. ❖ Need based drenching of (Metalaxyl + Mancozeb) or (Cymoxanil + Mancozeb) @400gm per acre is advised to control stem rot & collar rot.
Sugarcane	Vegetative stage	<ul style="list-style-type: none"> ❖ To control early shot borer in sugarcane foliar spray of Fipronil 5SC @400 ml/acre is advised.
Greengram	Flowering/ Fruiting	<ul style="list-style-type: none"> ❖ In summer greengram for the management of <i>Spodoptera litura</i> & pod borer spraying of Emamectin benzoate @80g per acre or prophenophos @400 ml per acre is recommended.

Vegetables	Flowering/ Fruiting	<ul style="list-style-type: none"> ❖ Since rain is anticipated, provide drainage in brinjal field. ❖ To control Fruit and shoot borer in Brinjal, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45 SC @70ml/acre or Emamectin benzoate @80g & Thiachlopid@250ml/l of water. ❖ To prevent from wilting or partial wilting in brinjal and tomato apply Neem or Karanjaa cake and spray Tebuconazole fungicide @ 1ml/liter to plant and plant base. ❖ Provide bamboo or wooden stake for better fruit set and growth of summer Tomato plants. ❖ Spraying Of Acephate @250g/acre along with neem oil formulation 600ml/acre to minimize damage caused by Red Pumpkin Beetle & Epilacna beetle in cucurbits. ❖ To control the little leaf in bitter guard & other viral diseases of cucurbits up root the infected plants & spray Flonicamid @60g/acre is recommended. ❖ Make the field clean and remove the weeds.
Fruit	Flowering /Fruiting	<p>Mango-</p> <ul style="list-style-type: none"> ❖ Spraying of NAA@1ml/4.5l is recommended to prevent the fruit drop & increase the fruit setting in mango. ❖ If infestation of Mealy bug & Mango hopper incidence in mango observed ETL, to control these pests, Spraying of Thiamethoxam 0.4 g/l of water or Lambda Cyhalothrin 4.9CS @1.25 ml/l of water recommended.
Livestock		<p>Light to Moderate rain is forecasted in the coming week, farmers are advised to keep livestock inside sheds, to prevent from rain water drenching. To prevent mortality due to thunder strike, grazing should be prevented.</p>

Mrs. Meera Mahanty
Jr. Agrometeorologist
AMFU, RRTTS, Ranital

Dr. Umasankar Nayak
ADR & NODAL OFFICER
AMFU, RRTTS, Ranital

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for IMD Weather forecast & Agromet advisory and “Damini” App for Lightning warning. For better agricultural benefits use.

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)