

Gramin Krishi Mausam Sewa India Meteorological Department Orissa University of Agriculture and Technology, Regional Research & Technology Transfer Sub Station,



Kirei, Sundergarh-770073

Dr. Sajeeb Kumar Biswasi, Senior Scientist & Nodal Officer, GKMS No-26(Eng.Bulletin) Dt -28.03.2024

Week No -13

District - Deogarh (North Western Plateau Agroclimatic Zone)

The mean maximum daily temperature was 35.0°C & mean minimum daily temperature was 21.6°C in Sundergarh district during last week. The District received in average 0.0 mm rainfall during the last week. Early sown rabi crops like Pulses, Oilseeds crop are at harvesting stage. The Vegetables are at harvesting stage. The overall crop condition is normal. The rabi paddy is in transplanted stage.

Forecast (Up to 02.04.2024)

Given by Met.Centre, IMD, Bhubaneswar

DISTRICT: DEOGARH- As per IMD Forecast, the district is may likely receive light rainfall for the next five days. The sky condition may be mainly clear to generally cloudy for the next five days. The wind speed is likely to remain within 7-10km/h. The Maximum Temperature and Minimum Temperature will likely to remain between 35-37°C and 24-25°C respectively for the next five days.

WARNING: There are chance of **light rainfall** at Deogarh district for coming 5days.

DISTRICT					
Date	29/03/2024	30/03/2024	31/03/2024	01/04/2024	02/04/2024
Rainfall (mm)	0	0	0	4	0
T.MAX(°C)	35	35	36	36	37
T.MIN (°C)	24	24	25	25	25
Cloud Cover	2	2	5	4	1
Rh Max (%)	44	58	53	40	36
Rh Min (%)	14	14	14	14	14
Wind speed	7	7	7	8	10
(kmph)					
Wind Direction	288	153	135	248	288
(deg)					

For further information, contact the Met. Centre, Aerodrom Area, IMD, Bhubaneswar, Tel.# 0674-2596116

Agromet Advisory

DISTRICT-DEOGARH.

General Advisory: In case of shortage of water, go for alternative row irrigation method for better utilization of water which saves around 50% water. Also the drip irrigation system in Orchard saves water. Sprinkling irrigation method reduces the effect of high temperature.

Crop specifi	c Advisory :	
Name of	Stage of the	Agromet Advisory
the Crop	Crop	
Sunmmer	Tillering stage	Due to rise in temperature, the BPH attacke may be seen, to manage this,
Paddy		make alleys in eeach meter gap for effective sunlight at the base of the plant,
		chemically spray in above ETL Ethofenfrox @ 1 ml per liter of water.
SRI Paddy	Tillering stage	Due to rise in temperature, provide irrigation when thin hair like carakes
		forms in the field. Run cono weeder to control the weeds.
Summer	Branching	Due to weather survey the infestation of YMV disease and to manage this
Pulses		apply Ridomil MZ @ 2g/liter of water.
Maize	Knee Height	Due to rise in temperature, provide irrigation at critical stages of Knee height
		stage
		To manage the Stem borer release T. Chilonis card @ 20,000/ac and repeat it after 10 days of first release.

Summer Groundnut	Peg Formation	Provide irrigation due to rise in temperature for easing Peg penetration and pod development After flowering, do not start inter cultivation work again, because from this time, the flower buds and its pods penetrate and set fruit.
Ragi	Tillering stage	Due to diurnal variation in temperature, After 3-4 weeks of planting finger millet, apply 12 kg of molasses per acre.
Animal		Due to rise in temperature provide the animal clean and sufficient drinking water.
Vegetables	Vegetative growth	Due to temperature effect in brinjal Foot & Shoot borer can be managed by application of Neem based pesticides 330 ppm @ 1 liter per acre.
Pisciculture	Growth stage	Due to rise in temperature, survey for oxygen deficit may be there.
Onion	Harvesting stage	Harvest the onion laving 2-3 cm from the neck and cut all the roots and dry it under shed during high temperature.

For any problem related to agriculture sector, please dial the toll free number 1551.

For better agricultural benefits use, **OUAT KALINGA** Products (Seedlings, Saapplings and QPM, Tissue culture plants, Fruits and vegetables, value added products, Fingerlings/Yearlings, Poultry chicks, Breeds, Mushrooma and spwn, Biofertilizer, Biopesticides, Vermi & Vermicompost and Farm implements)

IMD weather forecast & Agrometeorological Advisory of Odisha location is now available on Meghdoot mobile app in English & Odia Language.

Download: (Andriod: https://play.google.com/store/apps/details?id=com.aas.meghdoot)

ବି : ଦୃଷ୍ଟବ୍ୟ :ଓୟୂଏଟି ଶାଖା ଚିପଲିମା ଓ ଗୟାରିପାଲି ବିହନ ଗବେଷଣା ଫାର୍ମ ଠାରେ ସ୍ପର୍ଶ ଧାନ ବିହନ ୨୦ କିଗ୍ରା ବୟା ପିଛା ଟ. ୭୧୨ ୪୦ . ପଇସା ଚାଷୀ ନିମିତ୍ତ ବିକ୍ରି ହଉେଅଛି ।

Nodal Officer