



# Gramin Krishi Mausam Sewa

## India Meteorological Department

### Odisha University of Agriculture and Technology

#### Bhubaneswar -751 003

**Dr.T.R Mohanty**  
Nodal Officer

Week No.10

No. –222 (Eng. Bulletin)  
Dt. – 07.03.2024

**District – Jagatsinghpur (East and South-Eastern Coastal Plain Agroclimatic Zone)**

The mean maximum daily temperature was 31.8°C and mean minimum daily temperature was 24.8°C of the Jagatsinghpur district during the last week. The district received no rainfall during the last seven days. Transplanting of summer paddy is going to be completed. Sowing of summer pulses, oilseed and vegetables are also under progress. Mustard, Til and Groundnut are in harvesting stage. Harvesting of rabi paira pulses like Mung, Biri and Kulthi are continuing. Planting of Sugarcane is under progress. Crop coverage till date is 93.5 % of the total programme area. Overall crop condition is Normal.

Forecast (Up to 12.03.2024)

Given by Met. Centre, IMD, Bhubaneswar

**DISTRICT: JAGATSINGHPUR** – As per the forecast received from IMD, Jagatsinghpur district is likely to receive light rainfall on Friday & Saturday and dry weather on other days. The sky condition may remain clear to generally cloudy for the next five days. In the next 5 days, the wind speed will remain within 11-15 km/h. The daily maximum and minimum temperature are likely to remain 31-34°C and 21-25°C respectively for next five days.

DISTRICT	JAGATSINGHPUR				
Date	08/03/2024	09/03/2024	10/03/2024	11/03/2024	12/03/2024
Rainfall (mm)	7	5	0	0	0
T-MAX (C )	31	32	32	33	34
T-MIN (C )	25	23	21	21	21
Cloud Cover	7	7	1	0	0
Rh Max (%)	84	85	66	79	86
Rh Min (%)	55	57	42	44	50
Wind speed (kmph)	12	11	12	11	15
Wind Direction (deg)	189	116	90	108	198

For further information, contact the Met. Centre, Aerodrom Area, IMD, Bhubaneswar, Tel. # 0674-2596116.

**Agromet Advisory**

- ❖ If infestation of Bakanae disease is noticed in Paddy, spray, Carbendazim 50WP @ 200 g per acre in 200 litres of water and repeat the spray at 7-10 days' interval.
- ❖ In-case of Leaf blast infestation, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 80 g/acre or Isoprothiolane 40EC @ 300 ml/acre in 200 litres of water. Repeat the spray at 7-10 days' interval.
- ❖ Apply urea @ 35 kg/acre as top dressing at 20-25 DAT (tillering stage) in Paddy.
- ❖ If pre-emergence herbicides were not applied in paddy, to control weeds spray Bispyribacsodium @ 120 ml/acre at 10-12 DAS or, 2-3 leaf stage of weeds or, Penoxsulum 1.02% + Cyhalofop-butyl 5.1% @ 800ml/acre at 15-20 DAS in 140 litres of water.
- ❖ For controlling stem and root borer in cashew remove minimum width of the bark (only the damaged or tunneled portions) retaining at least 50% of the bark circumference.
- ❖ Transplanting of vegetables should be done in afternoon hour to avoid high temperature.
- ❖ Harvest the mustard crop as soon as the pods turn yellowish-brown. The crop is liable to shattering, hence it should be harvested just before the pods open to avoid losses. The harvested crop should be stacked in threshing floor for 5- days before threshing. After threshing dry the seeds in sunlight for few days to bring the moisture content of seeds up to 8 %.
- ❖ Harvest the potato crop which are at 75-85 days after planting and when majority of the leaves turn yellowish brown. For harvesting stop the irrigation 15 days before. All the harvested tubers should be kept in shed for 24 hours. Remove all the damaged and rotten tubers.
- ❖ To manage Tobacco Caterpillar pest in Groundnut, Blackgram, Greengram during primary stages of pest infestation proper surveillance should be done to collect and destroy egg mass and caterpillar of the pest. Monitor the emergence of adult moths by setting up Pheromone traps having Spodolure @ 6-8 nos/acre. Check the migration of caterpillars by dusting with Chlorpyrifos 1.5 % D @ 10 kg around the field.

**PADDY (Summer): Yellow stem borer management**

Due to rise in temperature there are chances of increase in stem borer infestation in paddy crop. To manage stem borer in paddy, broadcast granular insecticide Chlorantraniliprole 0.4 % GR @ 4-kg/acre or Cartap hydrochloride 4G @ 8-kg acre by mixing it with sand at 1:1 ratio or spray Chlorantraniliprole 18.5 % SC @ 60-ml/acre.

**SUGARCANE: Fertilizer Management**

Top dressing with Urea @ 60-kg per acre each at 45, 75 and 105 days after planting should be done. Apply the rest 20-kg of MOP at the time of the third top dressing with Urea. Do not top apply Urea after 120 days after planting as late application delayed maturity, reduces sucrose content and the total cane yield.

**MUSTARD: Downy Mildew disease**

To manage Downey Mildew disease in mustard crop spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre or Cymoxanil 8% + Mancozeb 64% W.P @500-gram/acre or Fosetyl-Al 80% WP @ 600- gram/acre.

**BRINJAL: Leaf hopper**

To manage leaf hopper in brinjal spray Thiamethoxam 25 % WG @ 40-gram/acre or Flonicamid 50% WG @ 60-gram/acre or Dinotefuran 20 % SG @ 80-gram/acre.

**COLE CROPS: Tobacco caterpillar management**

To manage tobacco caterpillar pest, regular monitoring of pest infestation should be carried out. Collect and destroy the caterpillars and egg masses in the early stages of pest attack. Spray Neem based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre by mixing in 200-litre of water. To manage these pests chemically spray Tolfenpyrad 15 % EC@ 400-ml/acre or Flubendiamide 20 % WG @ 20-gram/acre or Chlorfenapyr 10 % SC @ 400-ml/acre or Spinosad 45 % SC @ 75-ml/acre by mixing in 200-litre of water.

**PRINCIPAL NODAL OFFICER**

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

**IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.**  
Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>) (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)