



GRAMIN KRISHI MAUSAM SEWA
India Meteorological Department
Orissa University of Agriculture and Technology
Regional Research & Technology Transfer Station (RRTTS)
Bhawanipatna, Kalahandi – 766001



Bulletin No.19/2024 **WEEK NO. 10** **Monday/Dt.04.03.2024**

Weather Conditions recorded in last week at AMFU Bhawanipatna:

Weekly Rainfall (mm) and rainy days	1.8/00
Cumulative Rainfall (mm) and rainy day (w.e.f. 01.01.2024)	13.6/01
Mean weekly maximum temperature (°C)	34.3
Mean weekly minimum temperature (°C)	19.7

Weather Forecast from 05.03.2024 to 09.03.2024 for Kalahandi District:

DISTRICT	Date	Rainfall (mm)	T-MAX (°C)	T-MIN (°C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
KALAHANDI	05-03-2024	0	36	23	2	55	24	7	234
	06-03-2024	0	36	22	3	48	16	6	158
	07-03-2024	0	35	21	4	52	23	7	202
	08-03-2024	0	35	20	2	44	17	8	238
	09-03-2024	0	35	20	0	42	16	7	180

For further information, contact the IMD, Bhubaneswar, Tel.#0674-2596116

Weather Summery/Alert:

According to IMD forecast dry weather is very likely to prevail in next five days. Sky may remain clear to partly cloudy and wind may blow from Southerly to South Westerly direction with the speed of 006 to 008 km/hr. The maximum and minimum temperature may remain 35 to 36 °C and 20 to 23 °C respectively. Maximum RH is expected to vary between 42 – 55 % and minimum RH between 16 – 24 %.

Warning: Nil

Agromet – Advisory Bulletin for Kalahandi District

General Advisory:

- As per forecast received from IMD, there will be no rainfall in next five days. So, farmers are advised to check soil moisture status of the crop field and irrigation should be provided if required.
- Farmers are advised apply plant protection chemicals as the weather will be remain dry in subsequent five days.
- Farmers are advised to do weeding and hoeing in the standing crops and vegetables to protect them from weeds and be vigilant on infestation of pest & diseases.
- Apply balanced and appropriate fertilizers according to crop’s growth stage followed by irrigation.
- Keeping in view of rising day temperature, keep livestock inside shade. If the shade is built by RCC roof, then spreading of straw over the roof is advisable along with provision of clean water during day time to keep them hydrated.
- Vaccinate animals against FMD, BQ, HS, PPR & Goat Pox in regular interval.

SMS Advisory	Farmers are advised to select improved varieties of Okra like Utakala Gaurav, Arka Anamika, Arka Abhaya, Kasi Pragati etc.
Crop-Specific Advisory:	
Crop (Stage)	Advisory
Summer Paddy (Tillering Stage)	<p>Weed Management:</p> <ul style="list-style-type: none"> ▪ If pre-emergence herbicide not applied then, apply post emergence herbicide Bispyribac sodium @ 8 ml by mixing with 10 liters of water at 10-15 DAT or Penoxsulum 1.02% + cyhalofop-butyl 5.1% OD @ 4ml/ltr of water at 15-20 DAT for broad spectrum weed control. <p>Nutrient Management:</p> <ul style="list-style-type: none"> ▪ Apply urea 35kg/ac as first top dressing at tillering stage (20-25 DAT). <p>Disease and Insect Management:</p> <ul style="list-style-type: none"> ▪ If infestation of blast disease is noticed in direct seeded rice, spray Carbendazim 50 WP @ 1g/ ltr of water or Trifloxystrobin 25% + Tebuconazole 50% @ 4 g /10 ltrs of water or Edifenphos 50 EC @ 1 ml/ltr of water. ▪ In bacterial blight endemic areas, dip the seedling roots in 0.1% Plantomycin solution before transplanting for 30 minutes. ▪ Install 3 pheromone traps with 5mg lure/acre in the rice field for monitoring of the yellow stem borer and leaf folder just after the transplanting. Whenever the number of male moths/trap reaches 4 or 5, spray Azadiractin 0.15% neem seed kernel-based EC formulation @ 4 ml/ltr of water or



GRAMIN KRISHI MAUSAM SEWA
India Meteorological Department
Orissa University of Agriculture and Technology
Regional Research & Technology Transfer Station (RRTTS)
Bhawanipatna, Kalahandi – 766001



<p style="text-align: center;">Green gram/Black gram (Flowering/Fruiting)</p>	<p>apply Cartap hydrochloride 4G @ 10 kg/acre mixing with sand at 1:1 ratio.</p> <ul style="list-style-type: none"> Water management: Now the crop is at flowering stage. So flowering and grain filling stages are critical for green and black gram and water stress must be avoided during this period. <p>Disease and Insect Management:</p> <ul style="list-style-type: none"> Yellow Mosaic Virus/Jassids/Aphid/Thrips: If infection of YMV is noticed, then infected plants should be up rooted and buried in the soil. Installation of yellow sticky trap @ 8/acre to control sucking pests complex and blue sticky trap @ 8/acre for monitoring thrips at vegetative stage is recommended. Spray Thiomethoxam 25 WG @ 1g/ 5 ltr or Imidacloprid 17.8% SL @ 1.5 ml/ 5 ltr of water to control by chemical method. Powdery mildew: Spray Carbendazim 50% WP @1g/ltr or Penconazole 10% EC @ 0.5ml/ltr of water.
<p style="text-align: center;">Groundnut (Vegetative/Flowering)</p>	<ul style="list-style-type: none"> Water management: Flowering and pod development stages are critical for groundnut and water stress must be avoided at these stages. <p>Disease & Insect Management:</p> <ul style="list-style-type: none"> Aphid/Jassids/White fly/Thrips: Install yellow sticky trap @ 8/acre to monitor sucking pests complex and blue sticky trap @ 8/acre for monitoring thrips at vegetative stage. Tobacco Caterpillar: To monitor effectively install pheromone trap (Spodo lure) @ 8 no./ac at vegetative stage and replace the lure after 3 weeks.
<p style="text-align: center;">Hybrid Maize (Vegetative/Flowering/ Fruiting)</p>	<p>Insect Management:</p> <ul style="list-style-type: none"> Aphid & Thrips: Spray Thiomethoxam 25 WG @ 1g/ 5 ltr or Imidacloprid 17.8% SL @ 1.5 ml/ 5 ltr of water. Stem Borer: For effective control of stem borer spray Fipronil 5% SC or Profenophos 50% EC @ 2ml/ltr of water. FAW: Apply Emamectin Benzoate 5% SG @ 2 ml/ 5 ltr of water to control FAW infestation.
<p style="text-align: center;">Tomato (Flowering/Fruiting)</p>	<p>Disease and Insect Management:</p> <ul style="list-style-type: none"> Fruit & Shoot borer: Install pheromone trap @ 2nos/acre and for chemical control, spray Chlorantraniliprole 18.5% SC @ 3ml or Cytraniliprole 10.26% OD @ 18 ml or Indoxacarb 14.5% SC @ 10 ml by mixing with 10 liters of water. Leaf spot/Downy mildew/Powdery mildew: Spray Azoxystrobin 4.8% + Chlorothalonil 40% SC @ 6g /ltr of water. Leaf blight: Spray Copper oxychloride 50% WP @ 3g /ltr of water.
<p style="text-align: center;">Okra (Pre-sowing)</p>	<ul style="list-style-type: none"> Suitable time for sowing Okra @ 8-9 kg/ac during February and March. Apply well decomposed FYM @ 2 t/ac during land preparation. Seed treatment with Carbendazim 50% WP @ 2g/kg of seed should be done before sowing. Suitable Varieties: Utakala Gaurav, Arka Anamika, Arka Abhaya, Kasi Pragati etc.
<p style="text-align: center;">Chilli (Vegetative/Flowering/ Fruiting)</p>	<p>Disease and Insect Management:</p> <ul style="list-style-type: none"> Thrips: To manage thrips infestation install Blue sticky trap @ 8 nos./acre.For chemical control spray Emamectin Benzoate 5% SG @ 2 g or Fipronil 5% SC @ 10 ml or Imidacloprid 17.8% SL 1.5 ml by mixing with 5 ltrs of water. Fruit Borer: Install pheromone trap @ 8 nos./ac with Heli lure at vegetative stage and replace the lure after weeks.
<p style="text-align: center;">Brinjal (Vegetative/Flowering/ Fruiting)</p>	<p>Insect Management:</p> <ul style="list-style-type: none"> Fruit and Shoot borer: Install pheromone trap/Wata trap @ 8 nos. per acre area at vegetative stage and replace lure after 3 weeks. To control chemically, spray Spinosad 45% SC @ 5.5 ml by mixing with 15 liters of water or Thiodicarb 75 % WP @ 2ml/ltr of water. White fly: Install yellow sticky trap @ 8-10/acre or spray Neem Based Pesticide (Azadirachtin) 1500 PPM @ 3ml/ltr of water at initial stage of infestation.
<p style="text-align: center;">Onion (Vegetative)</p>	<p>Disease and Insect Management:</p> <ul style="list-style-type: none"> Purple blotch: Spray Difenconazole 25% EC or Tebuconazole 25.9% EC or Kitazin 48% EC @ 1g/ltr of water. Leaf blight: Spray Mancozeb 75% WP @ 3g/ltr of water. Thrips: Install blue sticky trap @ 8/ acre at vegetative stage. For chemical control, spray Thiomethoxam 25 WG @ 1g/ 5 liters of water or Acetamiprid 20% SP @ 1g/ 4 liters of water.



GRAMIN KRISHI MAUSAM SEWA
India Meteorological Department
Orissa University of Agriculture and Technology
Regional Research & Technology Transfer Station (RRTTS)
Bhawanipatna, Kalahandi – 766001



Livestock	
Cattle & Buffalo	<ul style="list-style-type: none">It is beneficial to add 5-10 % of fat in the diet of high yielding (10-15 ltr) cattle by reducing the fiber in the ration and add antioxidant like vitamin A & E in the diet. Sprinkle water over dairy cattle & buffalo twice or thrice a day to reduce their body temperature.
Goat & Sheep	<ul style="list-style-type: none">Allow the goats & sheeps to graze only from 7-11 AM in the morning & 4-7 PM in the evening.
Poultry	<ul style="list-style-type: none">Keep the curtains of poultry shed close during day time. If required, sprinkle water over them during day time for cooling effect.

Nodal Officer
GKMS, RRTTS, Bhawanipatna

N.B.: - Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for Agromet advisory and “Damini” App for Lightning warning. For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language. Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)