



GRAMIN KRISHI MAUSAM SEWA
India Meteorological Department
Orissa University of Agriculture and Technology
Regional Research & Technology Transfer Station (RRTTS)
Bhawanipatna, Kalahandi – 766001



Bulletin No.26/2024 **WEEK NO. 13** **Thursday/Dt.28.03.2024**

Weather Conditions recorded in last week at AMFU Bhawanipatna:

Weekly Rainfall (mm) and rainy days	0.2/010
Cumulative Rainfall (mm) and rainy day (w.e.f. 01.01.2024)	39.6/02
Mean weekly maximum temperature (°C)	36.4
Mean weekly minimum temperature (°C)	20.5

Weather Forecast from 29.03.2024 to 02.04.2024 for Kalahandi District:

DISTRICT	Date	Rainfall (mm)	T-MAX (°C)	T-MIN (°C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
KALAHANDI	29-03-2024	0	38	24	1	45	14	8	202
	30-03-2024	0	39	24	2	53	13	9	169
	31-03-2024	4	40	24	2	48	18	8	212
	01-04-2024	6	40	25	2	45	19	8	169
	02-04-2024	4	40	25	3	43	17	10	199

For further information, contact the IMD, Bhubaneswar, Tel.#0674-2596116

Weather Summery/Alert:

According to IMD forecast dry weather is very likely to prevail in next two days and light rainfall may occur in subsequent three days. Sky may remain mainly clear to partly cloudy and wind may blow from Southerly to South Westerly direction with the speed of 008 to 010 km/hr. The maximum and minimum temperature may remain 38 to 40 °C and 24 to 25 °C respectively. Maximum RH is expected to vary between 43 – 53 % and minimum RH between 13 – 19 %.

Warning: Nil

Agromet – Advisory Bulletin for Kalahandi District

General Advisory:

- Keeping in view of light rainfall during next five days, farmers are advised to harvest the matured crops.
- Considering light rainfall in coming week farmers are advised to apply plant protection chemicals and fertilizers according to the forecast.
- Regular monitoring of crop field is advisable for incidence of insects and diseases.
- Keep your crop field free from weed as it helps in spreading of insects and diseases. Farmers are advised to do mulching in vegetables to control weed population.
- As the day temperature is rising, keep livestock inside shade. If the shade is built by RCC roof, then spreading of straw over the roof is advisable along with provision of sufficient clean water during day time to keep them hydrated.

SMS Advisory	Install pheromone trap, sticky trap and solar light trap to control incidence of insects.
Crop and Livestock Specific Advisory:	
Crop (Stage)	Advisory
Summer Paddy (Panicle initiation)	<p>Nutrient Management:</p> <ul style="list-style-type: none"> Apply DAP @ 35kg/ac + MOP @ 11kg/ac at panicle initiation stage. <p>Disease and Insect Management:</p> <ul style="list-style-type: none"> Blast: If infestation of blast disease is noticed in direct seeded rice, spray Carbendazim 50 WP @ 1g/ ltr of water or Trifloxystrobin 25% + Tebuconazole 50% @ 4 g /10 ltrs of water or Edifenphos 50 EC @ 1 ml/ltr of water. Stem borer/Leaf folder: Apply nitrogenous fertilizer cautiously to avoid heavy infestation of stem borer. Collect and destroy egg masses of stem borer where ever possible. Install 3 pheromone traps with 5mg lure/acre in the rice field for monitoring of the yellow stem borer and leaf folder just after the transplanting. Whenever the number of male moths/trap reaches 4 or 5, spray Azadiractin 0.15% neem seed kernel-based EC formulation @ 4 ml/ltrs of water or apply Cartap hydrochloride 4G @ 10 kg/acre mixing with sand at 1:1 ratio or spray Chlorantraniliprole 18.5% SC @ 3ml or Flubendiamide 39.35% M/M SC @ 1ml by mixing with 10 liters of water.
Green gram/Black gram (Maturity/ Harvest)	<p>Post-Harvest Management:</p> <ul style="list-style-type: none"> Pods should be harvested at 70-80 % of the pods are matured and that should be sundried to bring the moisture content up to 8-10 percent of the seed. After harvesting of pods the plant should be mixed with the soil by ploughing, which enhances soil fertility.



GRAMIN KRISHI MAUSAM SEWA
India Meteorological Department
Orissa University of Agriculture and Technology
Regional Research & Technology Transfer Station (RRTTS)
Bhawanipatna, Kalahandi – 766001



Groundnut (Pod Development/Pod Maturity)	<p>Disease & Insect Management:</p> <ul style="list-style-type: none"> ▪ Leaf miner: Due to rise in day temperature infestation of leaf miner may be seen. Spray Profenophos 50% EC @ 2ml/liter of water if infestation is observed. ▪ Tobacco Caterpillar: To monitor effectively install pheromone trap (Spodo lure) @ 8 no./ac at vegetative stage and replace the lure after 3 weeks. ▪ Tikka: Spray Mancozeb 75% WP @ 3g/lit or Tebuconazole 29.5% EC @ 1.0 ml/lit or Carbendazim 50% WP @ 2g/ltr of water if tikka disease is noticed.
Hybrid Maize (Vegetative/Flowering/Fruiting)	<p>Insect Management:</p> <ul style="list-style-type: none"> ▪ Stem Borer: For effective control of stem borer spray Fipronil 5% SC or Profenophos 50% EC @ 2ml/ltr of water. ▪ FAW: Apply Emamectin Benzoate 5% SG @ 2 ml/ 5 ltr of water to control FAW infestation.
Tomato (Flowering/Fruiting)	<p>Disease and Insect Management:</p> <ul style="list-style-type: none"> ▪ Fruit & Shoot borer: Spray Chlorantraniliprole 18.5% SC @ 3ml or Cyntraniliprole 10.26% OD @ 18 ml or Indoxacarb 14.5% SC @ 10 ml by mixing with 10 liters of water. ▪ Leaf blight: Spray Copper oxychloride 50% WP @ 3g /ltr of water.
Chilli (Vegetative/Flowering/Fruiting)	<p>Disease and Insect Management:</p> <ul style="list-style-type: none"> ▪ Thrips/ Leaf curl virus: To manage thrips infestation install Blue sticky trap @ 8 nos./acre. For chemical control spray Emamectin Benzoate 5% SG @ 2 g or Fipronil 5% SC @ 10 ml or Imidacloprid 17.8% SL 1.5 ml by mixing with 5 ltrs of water. ▪ Fruit Borer: Install pheromone trap @ 8 nos./ac with Heli lure at vegetative stage and replace the lure after weeks.
Brinjal (Vegetative/Flowering/Fruiting)	<p>Insect Management:</p> <ul style="list-style-type: none"> ▪ Fruit and Shoot borer: Install pheromone trap/Wata trap @ 8 nos. per acre area at vegetative stage and replace lure after 3 weeks. To control chemically, spray Spinosad 45% SC @ 5.5 ml by mixing with 15 liters of water or Thiodicarb 75 % WP @ 2ml/ltr of water.
Livestock	
Cattle & Buffalo	<ul style="list-style-type: none"> ▪ It is beneficial to add 5-10 % of fat in the diet of high yielding (10-15 ltr) cattle by reducing the fiber in the ration and add antioxidant like vitamin A & E in the diet. Sprinkle water over dairy cattle & buffalo twice or thrice a day to reduce their body temperature.
Goat & Sheep	<ul style="list-style-type: none"> ▪ Allow the goats & sheeps to graze only from 7-11 AM in the morning & 4-7 PM in the evening.
Poultry	<ul style="list-style-type: none"> ▪ Keep the curtains of poultry shed close during day time. If required, sprinkle water over them during day time for cooling effect.

Nodal Officer
GKMS, RRTTS, Bhawanipatna

N.B.: - Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for Agromet advisory and “Damini” App for Lightning warning. For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language. Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)