

Gramin Krishi Mausam Sewa India Meteorological Department, AMFU, Malkangiri Odisha University of Agriculture & Technology



Krishi Vigyan Kendra, Malkangiri Mundaguda, P.O.+Dist.- Malkangiri, Odisha- 764045

Week No.:11 AGRO ADVISORY BULLETIN Date: 12.03.2024

<u>District - Malkangiri (South Eastern Ghat Zone)</u>

Weather condition during last four days									
Date	08.03.2024	09.03.2024	10.03.2024	11.03.2024					
Rainfall (mm)	0.0	0.0	0.0	0.0					
T. Max. (°C)	36	36	37	37					
T. Min. (° C)	22	21	24	25					

Forecast up to 17.03.2024 (According to MC, IMD, Bhubaneswar)

- ➤ Chances of clear sky to partly cloudy sky during next five days.
- > During next five days, chances of very light rainfall only on 17 March.
- Maximum temperature is expected to be 37-40°C and minimum temperature is expected to be 22-25°C.

Maximum and minimum relative humidity is expected to be 59-68%, and 20-23%, respectively.

Date	Rainfall (mm)	T. Max.	T. Min.	Cloud cover	RH Max. (%)	RH Min. (%)	Wind speed (Km/hour)	Wind direction (°)
13-03-2024	0	38	25	0	68	20	6	202
14-03-2024	0	39	25	0	65	20	7	234
15-03-2024	0	40	25	4	59	22	6	240
16-03-2024	0	40	23	4	59	21	7	202
17-03-2024	2	37	22	1	63	23	7	189

Agromet Advisory

General:

- Apply well decomposed Farm Yard Manure/ vermicompost during last ploughing in the field.
- ➤ Before raising any crop go for collection of soil sample and test it in nearby soil testing laboratory.
- > Apply balanced fertilizers in all the crops as per soil test based recommendations mentioned in the soil health card.

Rice:

- ➤ Those crop are at 3 week stage apply urea @ 35 kg/acre.
- ➤ Due to rise in temperature, the attack of stem borer has aggravated in rice. Three pheromone traps/acre may be installed in the rice field, whenever the number of male moths/trap reaches 4-5/day, spray Chlorantraniliprole 18.5% SC @ 0.3 ml/litre of water or Fipronil 5% SC @ 2 ml/litre of water.
- ➤ Bacterial leaf blight may reported in rice field, spray Plantomycin @ 1 g + Copper oxychloride 50% WP @ 3 g/litre of water. If necessary repeat at 15 days later.

Groundnut:

➤ Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut, to control these spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Pulses:

➤ To control vector of YMV in green gram spray Imidacloprid 17.8% SL @ 0.3 ml/litre of water or Acetamiprid 20% SP @ 0.3 g/litre of water.

Vegetables:

- ➤ White fly infestation may occur at this stage of tomato, to control it spray Imidacloprid 17.8% SL @ 0.3 ml/litre of water or Thiomethaxome 25% WG @ 0.4 g/litre of water.
- To control fruit and shoot borer in brinjal spray Spinosad 45% SC @ 70 ml/acre and Thiodicarb 75% WP @ 250 g/acre at 15 days interval.

Mango:

➤ Chances of mango hopper infestation in present situation, to control it spray Buprofezin 25% SC @ 1-2 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water.

Spray Alpha Napthyl Acetic Acid 4.5% SL (Planofix) @ 4.5 ml in 10 litre of water when tender fruits are in pea size and go for second spray when fruits are in marble size.

Mushroom:

For Oyster mushroom cultivation use Niger sticks as substrate.

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language. Download: (Android: https://play.google.com/store/apps/details?id=com.aas.meghdoot), (iOS: https://apps.apple.com/in/app/meghdoot/id1474048155)

Note: Contact nearby KVK Scientist or State Agriculture Officer/ Horticulture Officer for necessary help and suggestion.

Dr. Parshuram Sial Nodal Officer, GKMS, Malkangiri