



Week No.:12

AGRO ADVISORY BULLETIN

Date: 19.03.2024

District Mollrongini	South Eastann	Chat Zana)
District - Malkangiri	South Lastern	Gliat Lone)

Weather condition during last four days								
Date	15.03.2024	16.03.2024	17.03.2024	18.03.2024				
Rainfall (mm)	0.0	0.0	0.0	0.0				
T. Max. (°C)	37	38	36	31				
T. Min. (°C)	24	19	23	23				

Forecast up to 24.03.2024 (According to MC, IMD, Bhubaneswar)

- Chances of mainly clear to overcast sky during next five days.
- During next five days, chances of very light to light rainfall during 20 to 22 march and no rainfall during other days.
- Maximum temperature is expected to be 29-35°C and minimum temperature is expected to be 20-24°C.
- Maximum and minimum relative humidity is expected to be 62-76%, and 14-33%, respectively

Date	Rainfall (mm)	T. Max. (°C)	T. Min. (°C)	Cloud cover	RH Max. (%)	RH Min. (%)	Wind speed (Km/hour)	Wind direction (°)
20-03-2024	15	29	21	8	76	24	8	248
21-03-2024	5	29	20	8	72	33	6	112
22-03-2024	1	31	22	1	70	14	5	163
23-03-2024	0	33	23	1	63	14	6	207
24-03-2024	0	35	24	1	62	14	7	225

General:

Agromet Advisory

- > Apply well decomposed Farm Yard Manure/ vermicompost during last ploughing in the field.
- Before raising any crop go for collection of soil sample and test it in nearby soil testing laboratory.
- Apply balanced fertilizers in all the crops as per soil test based recommendations mentioned in the soil health card.

Rice:

- Due to rise in temperature, the attack of stem borer has aggravated in rice. Three pheromone traps/acre may be installed in the rice field, whenever the number of male moths/trap reaches 4-5/day, spray Chlorantraniliprole 18.5% SC @ 0.3 ml/litre of water or Fipronil 5% SC @ 2 ml/litre of water.
- Bacterial leaf blight may reported in rice field, spray Plantomycin @ 1 g + Copper oxychloride 50% WP @ 3 g/litre of water. If necessary repeat at 15 days later.

Groundnut:

Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut, to control these spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Pulses:

To control vector of YMV in green gram spray Imidacloprid 17.8% SL @ 0.3 ml/litre of water or Acetamiprid 20% SP @ 0.3 g/litre of water.

Onion

Prepare main field to transplant onion seedlings at eight week stage. Apply FYM 2ton, DAP 43kg, urea 27kg, MOP 33kg and Sulphur 20kg for one acre area transplanting.

Vegetables:

- ➤ White fly infestation may occur at this stage of tomato, to control it spray Imidacloprid 17.8% SL @ 0.3 ml/litre of water or Thiomethaxome 25% WG @ 0.4 g/litre of water.
- To control fruit and shoot borer in brinjal spray Spinosad 45% SC @ 70 ml/acre and Thiodicarb 75% WP @ 250 g/acre at 15 days interval.

Mango:

- Chances of mango hopper infestation in present situation, to control it spray Buprofezin 25% SC @ 1-2 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water.
- Spray Alpha Napthyl Acetic Acid 4.5% SL (Planofix) @ 4.5 ml in 10 litre of water when tender fruits are in pea size and go for second spray when fruits are in marble size.

For better agricultural benefits use, OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language. Download :(Android: https://play.google.com/store/apps/details?id=com.aas.meghdoot), (iOS: <u>https://apps.apple.com/in/app/meghdoot/id1474048155</u>)

Note: Contact nearby KVK Scientist or State Agriculture Officer/ Horticulture Officer for necessary help and suggestion.

Dr. Parshuram Sial Nodal Officer, GKMS, Malkangiri