



No.26 (English Bulletin)

Week no-13

Date: 28.03.2024

West Central Table Land Zone

Sambalpur

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is a chance of occurrence of very light rainfall along with partly cloudy sky is expected at the middle of upcoming week. The maximum and minimum temperature is likely to be ranges between 38-39°C and 24-25 respectively during this week. The maximum and minimum relative humidity may vary between 33-46% and 13-15% respectively. The S/SW/NW with a speed of 7-11 km/hr is predicted during upcoming days.

| Date | 29-03-2024 | 30-03-2024 | 31-03-2024 | 01-04-2024 | 02-04-2024 |
|---------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 0 | 0 | 2 | 3 | 0 |
| Max Temperature (°C) | 38 | 38 | 39 | 39 | 39 |
| Min Temperature (°C) | 24 | 24 | 25 | 25 | 25 |
| Total cloud cover (okta) | 2 | 2 | 4 | 4 | 1 |
| Max Relative Humidity (%) | 33 | 46 | 40 | 42 | 37 |
| Min Relative Humidity (%) | 14 | 13 | 14 | 14 | 15 |
| Wind speed (km/h) | 7 | 7 | 9 | 8 | 11 |
| Wind direction (deg) | 180 | 255 | 236 | 297 | 285 |

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

General Advisory

- As there is a chance of dry weather condition along with increase maximum temperature is expected during upcoming week so farmers are advised to apply need based irrigation in crop field and conduct fertilizer application and spraying operations during that period. But due to chance of occurrence of cloudy weather along with very light rain at the middle of upcoming week farmers are advised to take proper care during spraying operations and keep the harvested produce in safe place. Mulching should be done with crop residue, straw /polythene to conserve soil moisture as well as maintain soil temperature.
- Present weather condition is favourable for increasing the population of different diseases and insects in crop field so its regular monitoring should be advised.
- Avoid grazing of animal during noon hours in open place because there is a chance of increase in maximum temperature in upcoming days.

Crop and livestock Specific Agromet Advisory

| Name of the crop | Crop Stage | Agromet Advisories |
|----------------------------|-------------------------------------|--|
| Rice | Tillering | <ul style="list-style-type: none"> Farmers are advised to apply top dressing of urea fertilizer in rice plant in tillering stage under adequate moisture condition. To control yellow stem borer, spraying of Chlorantraniliprole 18.5% SC @ 0.3ml or Fipronil 5% @ 2.5ml per liter of water should be advised under clear sky. For BLB control, farmer should spray Plantomycin @ 1g/liter of water using 200 liters of water per acre or Streptocycline (150 mg) + Copper oxychloride 1g/litre of water twice at an interval of 8 days. |
| Green Gram (Summer) | Vegetative / flowering stage | <ul style="list-style-type: none"> To control aphids attack in pulse crop, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water and for yellow mosaic virus (transmitted through white fly) spraying of Acetamiprid 20% SP @ 40 g/acre in 200-250 liters of water should be advised under clear weather. Yellow sticky traps or cards @ 10/ac. may reduce the whitefly population. |

| | | |
|------------------------|---|--|
| Maize | Vegetative stage | <ul style="list-style-type: none"> • Farmers should apply top dressing of nitrogen fertilizer under proper soil moisture condition at this stage. • To control stem borer in Maize field, spraying of Fipronil 5% SC or Profenophos 50% EC @ 2ml and for fall army worm, Emamectin Benzoate @ 0.4 g per liter of water should be advised. |
| Mango | Fruiting stages | <ul style="list-style-type: none"> • To control mango hopper in Mango tree, spraying of insecticides like Acephate (1g/L) or Chloropyriphos (0.5 ml/L) or Thiomethoxam (0.3-0.4 g/L) and for powdery mildew disease, spraying of soluble sulphur (2 g/ L) should be advised under clear weather condition. • Spraying of Planofix @ 1ml /4.5 liters of water should be advised at Mango with pea size fruit stage to reduce fruit drops. • Farmers shouldn't apply hormone and insecticide at same time. There should be at least one week gap between their applications. |
| Vegetable crops | Vegetative stage / flowering/ Fruiting | <ul style="list-style-type: none"> • Proper monitoring against root and fruit rot disease infestation in vegetable crops should be advised. • To control leaf eating caterpillar and fruit borer / shoot and fruit borer in Tomato, Brinjal and Okra spraying of Emamectin Benzoate @ 0.4 g per liter of water should be advised under clear weather. • Early (Alternaria) blight infection is observed in some tomato fields so spraying of Tebuconazole 50% + Trifloxystrobin 25% @ 0.4g per liter water should be advised. • In vegetative stage of tomato, leaf minor and white fly attack observed in some farmers field so application of NSKE 5% with a sticker found to be effective. Leaf minor population may boost up due to excess use of nitrogen fertilizer so its judicious use should be advised. s • Spraying of Tebuconazole 50% + Trifloxystrobin 25% @ 0.4g per liter water should be advised to control purple blotch disease in Onion and in case of thrips attack, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water or Acetamiprid 20% SP @ 1g per 4 liters of water should be advised. |
| Cow/Buffalo | | <ul style="list-style-type: none"> • To protect the animal from heat, farmers should make provision to provide fresh clean and cold drinking water to the animal along with salt because temperature is expected to be increased in upcoming days. Keep animals in shade and don't make them work between 11 am to 4 pm. Cover the shed roof with straw, paint it white or plaster with dung-mud to reduce temperature. |

| | |
|---|--|
| <p>Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima</p> | <p>Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima</p> |
|---|--|

For better agricultural benefits, use **OUAT KALINGA products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)**

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile**

app(<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information.