



**GraminKrishiMausamSewa**  
**India Meteorological Department**  
**Orissa University of Agriculture and Technology**  
**RRTTS, Chiplima, Sambalpur**



No.29 (English Bulletin)

Week no-15

Date: 09.04.2024

**West Central Table Land Zone**

**Bargarh**

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is a chance of occurrence of light rainfall along with partly to generally cloudy sky is expected in upcoming two days and thereafter very less chance of rainfall along with partly clear or partly cloudy sky is forecasted. The maximum and minimum temperature is likely to be ranges between 27-37<sup>0</sup>C and 23-26<sup>0</sup>C respectively during this week. The maximum and minimum relative humidity may vary between 44-62% and 18-31% respectively. The S/SW/SE wind with a speed of 6-11 km/hr is predicted during upcoming days.

Date	10-04-2024	11-04-2024	12-04-2024	13-04-2024	14-04-2024
Rainfall (mm)	10	2	0	0	0
Max Temperature (° C)	27	32	35	37	37
Min Temperature (° C)	23	24	26	26	26
Total cloud cover (okta)	7	5	4	3	2
Max Relative Humidity (%)	58	62	55	44	46
Min Relative Humidity (%)	18	30	31	21	23
Wind speed (km/h)	11	10	8	6	9
Wind direction (deg)	180	198	155	180	252

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

**General Advisory**

- As per ERFs model of IMD, rainfall is above normal, maximum and minimum temperature is below normal and normal respectively for Odisha during 14.04.2024 to 20.04.2024 and there is chance of occurrence of light rainfall in upcoming one or two days, so farmers should conduct spraying operations and schedule the irrigation accordingly.

**Crop and livestock Specific Agromet Advisory**

Name of the crop	Crop Stage	Agromet Advisories
<b>Rice</b>	<b>Panicle initiation / 50% flowering</b>	<ul style="list-style-type: none"> <li>Farmers are advised to maintain adequate moisture in paddy field at this stage and apply final top dressing of nitrogen fertilizer after draining out excess water from field.</li> <li>To control yellow stem borer, spraying of Chlorantraniliprole 18.5% SC @ 0.3ml or Fipronil 5% @ 2.5ml per liter of water should be advised under clear sky.</li> <li>For BLB control, farmer should spray Plantomycin @ 1g/liter of water using 200 liters of water per acre or Streptocycline (150 mg) + Copper oxychloride 1g/litre of water twice at an interval of 8 days.</li> </ul>
<b>Green Gram (Summer)</b>	<b>flowering / Pod formation stage</b>	<ul style="list-style-type: none"> <li>Looking to the previous rainfall condition, farmers are advised to drain out excess water from crop field.</li> <li>To control aphids attack in pulse crop, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water should be advised under clear weather.</li> <li>For pod borer attack, spraying of Chlorantraniliprole 18.5% SC @ 0.3ml or Emamectin Benzoate @ 0.4 g or Prophenofos @ 2ml per liter of water should be advised when sky is clear.</li> </ul>
<b>Maize</b>	<b>Vegetative/ Tasseling</b>	<ul style="list-style-type: none"> <li>Farmers should apply top dressing of nitrogen fertilizer under proper soil moisture condition at this stage.</li> </ul>

	<b>stage</b>	<ul style="list-style-type: none"> <li>To control stem borer in Maize field, spraying of Fipronil 5% SC or Profenophos 50% EC @ 2ml and for fall army worm, Emamectin Benzoate @ 0.4 g per liter of water should be advised.</li> </ul>
<b>Mango</b>	<b>Fruiting stages</b>	<ul style="list-style-type: none"> <li>To control mango hopper in Mango tree, spraying of insecticides like Acephate (1g/L) or Chloropyriphos (0.5 ml/L) or Thiomethoxam (0.3-0.4 g/L) and for powdery mildew disease, spraying of soluble sulphur (2 g/ L) should be advised under clear weather condition.</li> <li>Spraying of Planofix @ 1ml /4.5 liters of water should be advised at Mango with pea size fruit stage to reduce fruit drops.</li> <li>Farmers shouldn't apply hormone and insecticide at same time. There should be at least one week gap between their applications.</li> <li>Water logged condition should be avoided in Mango orchard, for this proper drainage of excess water from field should be recommended.</li> </ul>
<b>Vegetable crops</b>	<b>Vegetative stage / flowering/ Fruiting</b>	<ul style="list-style-type: none"> <li>Looking to the previous rainfall condition, farmers are advised to drain out excess water from vegetable field to avoid water logged condition which may leads to root rot and fruit rot diseases.</li> <li>To control leaf eating caterpillar and fruit borer / shoot and fruit borer in <b>Tomato, Brinjal and Okra</b> spraying of Emamectin Benzoate @ 0.4 g per liter of water should be advised under clear weather.</li> <li>Spraying of Tebuconazole 50% + Trifloxystrobin 25% @ 0.4g per liter water should be advised to control <b>purple blotch</b> disease in <b>Onion</b> and in case of thrips attack, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water or Acetamiprid 20% SP @ 1g per 4 liters of water should be advised.</li> </ul>
<b>Cow/Buffalo</b>		<ul style="list-style-type: none"> <li>To protect the animal from heat, farmers should make provision to provide fresh clean and cold drinking water to the animal along with green grass, protein-fat bypass supplement, mineral mixture and salt because temperature is expected to be increased after some days. Keep animals in shade and don't make them work between 11 am to 4 pm. Cover the shed roof with straw, paint it white or plaster with dung-mud to reduce temperature. In extreme heat condition, spray water and take the animal to a water body to cool off. Make them graze during cooler hours.</li> </ul>

<b>Ms. Jubuli Sahu</b> <b>Junior Agrometeorologist,</b> <b>GKMS, RRTTS, Chiplima</b>	<b>Dr. Sanjukta Mohapatra</b> <b>ADR cum Nodal Officer</b> <b>RRTTS, Chiplima</b>
--	---

**For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)**

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information.