



West Central Table Land Zone

Boudh

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is very less chance of occurrence of rainfall along with mainly clear to partly cloudy sky is expected during upcoming week. The maximum and minimum temperature is likely to be ranges between 28-41^oC and 21-26^oC respectively during this week. The maximum and minimum relative humidity may vary between 37-47% and 15-27% respectively. The W/SW/SE wind with a speed of 6-8 km/hr is predicted during upcoming days.

Date	13-04-2024	14-04-2024	15-04-2024	16-04-2024	17-04-2024
Rainfall (mm)	0	0	0	0	0
Max Temperature (° C)	28	32	36	40	41
Min Temperature (° C)	21	22	24	26	24
Total cloud cover (okta)	4	3	1	1	2
Max Relative Humidity (%)	44	47	42	45	37
Min Relative Humidity (%)	27	18	20	18	15
Wind speed (km/h)	6	8	6	7	7
Wind direction (deg)	150	270	202	169	206

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

General Advisory

- There is a chance of increase in temperature after one day, so farmer should take precaution accordingly.
- Apply light and frequent irrigation to the standing crops, increase the frequency of irrigation at critical growth stages, irrigate only during the evening or early morning and use sprinkler irrigation if possible.
- Mulching should be done with crop residue, straw / polythene to conserve soil moisture as well as maintain soil temperature.
- Avoid grazing of animal during noon hours in open place because there is a chance of increase in maximum temperature in upcoming days.
- Past weather condition is favourable for increasing the population of various insects and diseases in both field as well as vegetable crops so its regular monitoring should be advised.

Crop and livestock Specific Agromet Advisory

Name of the crop	Crop Stage	Agromet Advisories
Rice	Panicle initiation / 50% flowering	<ul style="list-style-type: none"> • Farmers are advised to maintain adequate moisture in paddy field at this stage and apply final top dressing of nitrogen fertilizer after draining out excess water from field. • To control yellow stem borer, spraying of Chlorantraniliprole 18.5% SC @ 0.3ml or Fipronil 5% @ 2.5ml per liter of water should be advised under clear sky. • For BLB control, farmer should spray Plantomycin @ 1g/liter of water using 200 liters of water per acre or Streptocycline (150 mg) + Copper oxychloride 1g/litre of water twice at an interval of 8 days.
Green Gram (Summer)	flowering / Pod formation stage	<ul style="list-style-type: none"> • Looking to the previous rainfall condition, farmers are advised to drain out excess water from crop field. • To control aphids attack in pulse crop, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water should be advised under clear weather. • For pod borer attack, spraying of Chlorantraniliprole 18.5% SC @

		0.3ml or Emamectin Benzoate @ 0.4 g or Prophenofos @ 2ml per liter of water should be advised when sky is clear.
Maize	Vegetative/ Tasseling stage	<ul style="list-style-type: none"> • Farmers should apply top dressing of nitrogen fertilizer under proper soil moisture condition at this stage. • To control stem borer in Maize field, spraying of Fipronil 5% SC or Profenophos 50% EC @ 2ml and for fall army worm, Emamectin Benzoate @ 0.4 g per liter of water should be advised.
Groundnut	Peak flowering/ Peg and Pod formation	<ul style="list-style-type: none"> • Application of Gypsum @ 200 kg per acre should be advised at flowering stage on soil surface near to the base of the plant after occurrence of rainfall under adequate moisture condition so that calcium is picked up by the peg and the developing pods
Mango	Fruiting stages	<ul style="list-style-type: none"> • To control mango hopper in Mango tree, spraying of insecticides like Acephate (1g/L) or Chloropyriphos (0.5 ml/L) or Thiomethoxam (0.3-0.4 g/L) and for powdery mildew disease, spraying of soluble sulphur (2 g/ L) should be advised under clear weather condition. • Spraying of Planofix @ 1ml /4.5 liters of water should be advised at Mango with pea size fruit stage to reduce fruit drops. • Farmers shouldn't apply hormone and insecticide at same time. There should be at least one week gap between their applications. • Water logged condition should be avoided in Mango orchard, for this proper drainage of excess water from field should be recommended.
Vegetable crops	Vegetative stage / flowering/ Fruiting	<ul style="list-style-type: none"> • Looking to the previous rainfall condition, farmers are advised to drain out excess water from vegetable field to avoid water logged condition which may leads to root rot and fruit rot diseases. • To control leaf eating caterpillar and fruit borer / shoot and fruit borer in Tomato, Brinjal and Okra spraying of Emamectin Benzoate @ 0.4 g per liter of water should be advised under clear weather. • Spraying of Tebuconazole 50% + Trifloxystrobin 25% @ 0.4g per liter water should be advised to control purple blotch disease in Onion and in case of thrips attack, spraying of imidacloprid @ 0.3 ml or Thiamethoxam @ 0.3 g per liter of water or Acetamiprid 20% SP @ 1g per 4 liters of water should be advised.
Cow/Buffalo	<ul style="list-style-type: none"> • There is a chance of increase in temperature in upcoming week so to protect the animal from heat, farmers should make provision to provide fresh clean and cold drinking water to the animal along with green grass, protein-fat bypass supplement, mineral mixture and salt because temperature is expected to be increased after some days. Keep animals in shade and don't make them work between 11 am to 4 pm. Cover the shed roof with straw, paint it white or plaster with dung-mud to reduce temperature. In extreme heat condition, spray water and take the animal to a water body to cool off. Make them graze during cooler hours. 	

<p>Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima</p>	<p>Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima</p>
---	--

For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)

IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information.