



# Gramin Krishi Mausam Sewa

## India Meteorological Department

### Odisha University of Agriculture and Technology

#### Bhubaneswar -751 003

**Dr.T.R Mohanty**  
Nodal Officer

Week No.14

No. –363 (Eng. Bulletin)  
Dt. – 02.04.2024

**District – Jagatsinghpur (East and South-Eastern Coastal Plain Agroclimatic Zone)**

The mean maximum daily temperature was 32.8°C and mean minimum daily temperature was 27.0°C of the Jagatsinghpur district during the last week. The district received no rainfall during the last seven days. Transplanting of summer paddy is completed. Sowing of summer pulses, oilseed and vegetables are also under progress. Harvesting of rabi crops like Mustard, Groundnut, Mung, Biri and Kulthi are completed. Crop coverage till date is 99 % of the total programme area. Overall crop condition is Normal.

Forecast (Up to 07.04.2024)

Given by Met. Centre, IMD, Bhubaneswar

**DISTRICT: JAGATSINGHPUR** – As per the forecast received from IMD, dry weather may prevail over the district of Jagatsinghpur for the next five days. The sky condition may remain clear to mainly clear for the next five days. In the next 5 days, the wind speed will remain within 19-28 km/h. The daily maximum and minimum temperature are likely to remain 33-34°C and 28-29°C respectively for next five days.

DISTRICT	JAGATSINGHPUR				
Date	03/04/2024	04/04/2024	05/04/2024	06/04/2024	07/04/2024
Rainfall (mm)	0	0	0	0	0
T-MAX (C )	33	33	33	34	34
T-MIN (C )	28	28	29	29	29
Cloud Cover	0	1	0	0	0
Rh Max (%)	79	92	86	83	93
Rh Min (%)	45	49	60	48	43
Wind speed (kmph)	24	22	28	25	19
Wind Direction (deg)	207	202	202	204	197

For further information, contact the Met. Centre, Aerodrom Area, IMD, Bhubaneswar, Tel. # 0674-2596116.

**Agromet Advisory**

- ❖ In paddy crop, Panicle initiation, Panicle emergence, flowering and milking stages are the critical stages. In these stages water scarcity or deficit of water causes comparatively greater reduction in yields. Hence, water deficit during these stages should be avoided.
- ❖ Cover the rows with dry leaves in Sugarcane. Top-dress with 4% potassium to protect crop from water scarcity during summer.
- ❖ Apply sprinkler irrigation to bring down the heat effect. Conserve upto 50% water by irrigating alternate rows.
- ❖ The high temperature increases the water requirement of maize crop. Irrigate the crop. Never allow water to remain stagnant for more than six hour in the field.
- ❖ Spread polythene in irrigation channels to save water wastage during irrigation in sandy and loamy soil.
- ❖ High temperature in day hours may lead to wilting in tomato and brinjal. Apply 1g Streptocyclin in 10 liter water in root zone.
- ❖ Mulch coconut leaves 1.8 m around a coconut tree to reduce water loss.
- ❖ Cover with paper caps on the newly planted vegetable seedlings during day time.
- ❖ In shallow ponds, harvest fingerlings, if water temperature rises to 30 °C. Ensure enough water depth before introducing fingerlings in a pond.

**General Advisory**

**PADDY (Summer): Yellow stem borer management-** Due to rise in temperature there are chances of increase in stem borer infestation in paddy crop. To manage Yellow Stem Borer in paddy spray Fipronil 5%SC @ 400ml/acre or Flubendiamide 20% WG @ 50g/acre or Chlorantraniliprole 18.5% SC @ 60ml /acre or Chromafenozide 80%WP @ 50g/acre.

**Blast Disease Management** - To manage blast disease spray Hexaconazole 5 % SC @ 400-ml/acre or Azoxystrobin 18.2% + Difenconazole 11.4 % S.C @ 200-ml/acre or Tebuconazole 50%+ Trifloxystrobin 25 % WG @ 80-gram/acre.

**BLACK GRAM & GREEN GRAM: Tobacco Caterpillar Management**

Check the migration of caterpillars by dusting with Chlorpyriphos 1.5 % D @ 10 kg around the field. For chemical control spray Novaluron 5.25 % + Indoxacarb 4.25 % @ 400-ml/acre or Lambda Cyhalothrin 4.9 CS @ 300 ml or Flubendiamide 20 % WG @ 120 gram. For spraying of insecticides 200 litre of water per acre is required.

**SUGARCANE: Early Shoot Borer** :To manage early shoot borer in Sugarcane, irrigate the crop frequently during hot months. Timely cutting the attacked shoot at or just below the ground level ensures destruction of most of the caterpillar. Collect and destroy the egg masses manually. In furrow, apply Fipronil 0.3% GR @ 10 kg/acre at the time of first earthing up.

**BRINJAL: Brinjal shoot and fruit borer**

To manage this insect, install 20 nos. of pheromone trap having Leucin Lure per acre. During primary stage of pest infestation spray neem-based pesticide (Azadirachtin) 1500 PPM @ 600-ml/acre by mixing in 200-litre of water. Go for chemical control if there is 4% withering of terminal shoot or 14% fruits infested with borers. For chemical control spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/acre or Chlorantraniliprole 18.5 % SC @80-ml/acre or Spinetoram 11.7 % SC @ 200-ml /acre. Do not repeat the same insecticide. Use any of the above insecticides alternately at 15-days interval.

**OKRA: Whitefly Infestation:** To prevent whitefly infestation in okra seeds should be treated with Imidacloprid @ 7-gram/kg of seeds before sowing. Install 8-10 nos of yellow sticky traps starting from 2-3 leaf stage of the crop.

**ONION:** After harvesting of onion, cut the tops leaving 2.0-2.5 cm above bulb. The harvested bulb should be dried in well ventilated concrete floored room. Well dried bulbs can be stored for longer period.

**MANGO: Mango hopper** -To control mango hoppers spray Imidacloprid 17.8% SL @ 5 ml/15 litre of water or Thiamethoxam 25% W.G @ 3ml/15litre of water or Acetamiprid 20% S.P. @ 4g /15litre of water. Application of smoke below the plants can also help in reducing the pest population.

**PRINCIPAL NODAL OFFICER**

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

**IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.**

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>) (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)