



**Gramin Krishi Mausam Sewa**  
**India Meteorological Department, AMFU: Semiliguda**  
**Odisha University of Agriculture & Technology**  
**Regional Research & Technology Transfer Station**  
**Semiliguda, Post Box No: 10, Sunabeda, Koraput-763002**



**Week No.: 17**

**AGRO ADVISORY BULLETIN**

**Date: 23.04.2024**

**District - Nabarangpur (Eastern Ghat High Land Zone)**

**Forecast up to 28.04.2024 (According to MC, IMD, Bhubaneswar)**

- Chances of mainly clear to generally cloudy sky during next five days.
- Chances of light rainfall on 24<sup>th</sup> April & no rainfall during 25-28 April.
- Maximum temperature is expected to be 39-41°C and minimum temperature is expected to be 24-25°C.
- Maximum and minimum relative humidity is expected to be 40-43%, and 17-23%, respectively.

Date	Rainfall (mm)	T. Max. (°C)	T. Min. (°C)	Cloud cover	RH Max. (%)	RH Min. (%)	Wind speed (Km/hour)	Wind direction (°)
24-04-2024	4	39	24	6	42	23	9	180
25-04-2024	0	39	24	4	43	21	10	198
26-04-2024	0	41	25	2	42	17	10	287
27-04-2024	0	41	25	2	41	20	8	285
28-04-2024	0	41	25	2	40	20	8	255

**Agromet Advisory**

**General:**

- Plant protection measures may be taken up when weather becomes clean and dry, avoid in rainy days.
- Take up summer deep ploughing by using recent rainfall to destroy the weed seeds and eggs of insect pests.
- Farmers are advised for not to burn the residue of kharif crops in the fields because it not only pollute the environment but also causes the severe health problems.

**Rice:**

- At panicle initiation stage apply 18 kg urea in medium duration paddy and 13 kg urea in short duration paddy.
- If blast infestation found in rice, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g/litre of water in clear and dry weather..
- Due to rise in temperature, the attack of stem borer has aggravated in rice. To control it spray Chlorantraniliprole 18.5% SC @ 0.3 ml/litre of water or Fipronil 5% SC @ 2 ml/litre of water.
- Bacterial leaf blight may reported in rice field, spray Plantomycin @ 1 g + Copper oxychloride 50% WP @ 3 g/litre of water. If necessary repeat at 15 days later.

**Oilseeds:**

- Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut, to control these spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

**Onion:**

- Prepare main field to transplant onion seedlings at eight week stage. Apply FYM 2ton, DAP 43kg, urea 27kg, MOP 33kg and Sulphur 20kg for one acre area.

**Vegetables:**

- White fly infestation may occur at this stage of tomato, to control it spray Imidacloprid 17.8% SL @ 0.3 ml/litre of water or Thiomethaxome 25%WG @ 0.4 g/litre of water.
- If bacterial wilting observes in tomato, apply Plantomycin @ 1.5 g and Copper Oxychloride @ 3 g/litre of water.
- To control fruit and shoot borer in brinjal spray Spinosad 45% SC @ 70 ml/acre and Thiodicarb 75% WP @ 250 g/acre at 15 days interval.

For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & VermiCompost and Farm Implements).

**IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on Meghdoot mobile app in English and Odia language.**  
Download :( Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)

**Note: Contact nearby KVK Scientist or State Agriculture Officer/ Horticulture Officer for necessary help and suggestion.**

---

**Dr. Parshuram Sial**  
**Nodal Officer, GKMS, Semiliguda**