



**Gramin Krishi Mausam Sewa**  
**India Meteorological Department**  
**Orissa University of Agriculture and Technology,**  
**Regional Research & Technology Transfer Sub Station,**  
**Kirei, Sundergarh-770073**



**Dr. Sajeed Kumar Biswasi**

Senior Scientist & Nodal Officer, GKMS

No-34(Eng.Bulletin)

Dt -26.04.2024

**Week No - 17**

**District –Sundergarh (North Western Plateau Agroclimatic Zone)**

The mean maximum daily temperature was **40.6<sup>0</sup>C** & mean minimum daily temperature was **23.6<sup>0</sup>C** in Sundergarh district during last week. The District received in average **0.0 mm** rainfall during the last week. Early sown summer crops like Pulses, Oilseeds crop are at harvesting stage. The Vegetables are at harvesting stage. The overall crop condition is normal. The rabi paddy is in Panicle emergence to anthesis stage.

**Forecast (Up to 01.05.2024)**

**Given by Met.Centre, IMD, Bhubaneswar**

**DISTRICT: SUNDERGARH-** As per the forecast issued by India Meteorological Department, the district is likely receive no rainfall for the next 5 days. The sky condition is clear sky to mainly clear for the next five days .The wind speed is likely to remain within 8-11km/h. The Maximum Temperature and Minimum Temperature is forecasted to remain between 41-44<sup>0</sup>C and 23-25<sup>0</sup>C respectively for the next five days.

**WARNING :** The district is likely receive no rainfall in Sundergarh district for coming 5 days. .

DISTRICT	27/04/2024	28/04/2024	29/04/2024	30/04/2024	01/05/2024
Date	27/04/2024	28/04/2024	29/04/2024	30/04/2024	01/05/2024
Rainfall (mm)	0	0	0	0	0
T.MAX(°C)	43	44	44	43	41
T.MIN (°C)	25	25	25	24	23
Cloud Cover	1	2	2	0	1
Rh Max (%)	24	30	33	26	23
Rh Min (%)	9	9	12	8	9
Wind speed (kmph)	8	9	8	10	11
Wind Direction (deg)	248	244	270	287	287

For further information,contact the Met. Centre,Aerodrom Area, IMD, Bhubaneswar, Tel.# 0674-2596116 34

**DISTRICT: SUNDERGARH-Agromet Advisory**

**General Advisory :** In this case, provide irrigation to crops in critical stage specially during evening hours. In case of shortage of water, go for alternate row irrigation method for better utilization of water which saves around 50% water. Also the drip irrigation system in Orchard saves water. Sprinkling irrigation method reduces the effect of high temperature. Farmers are advised to avoid exposure outside during the pick heat hours of the day.

**Crop specific Advisory :**

Name of the Crop	Stage of the Crop	Agromet Advisory
Summer Paddy	PI stage	Due to rise in temperature, the BPH attack may be seen, to manage this, make alleys in every meter gap for effective sunlight at the base of the plant, chemically spray in case of above ETL Thiomethoxam @ 3 g per 15 liters of water. Manage to keep ponding water in the field. Provide critical stage irrigation to crop for reducing the effect of heat wave.
SRI Paddy	PI stage	Leaf folder may attack, go for spraying Lambdacyalothrin @ 1ml per liter of water.
Summer Pulses	Flowering stage	Due to weather survey the infestation of YMV disease and to manage this apply Ridomil MZ @ 2g/liter of water. Provide irrigation at flowering stage.

<b>Summer Groundnut</b>	<b>Pod Development</b>	Provide irrigation due to rise in temperature at pod development stage.
<b>Sunflower</b>	<b>Head Formation and seed setting</b>	For good seed filling, apply 2-3 g of Borax/liter of water. Early sown crop when the lower side of head turns yellow, go for cutting the head and dry it under sun for 4-5 days. If fruit borer attacks then go for spraying Bioinsecticide NPV @ 250 LE mixed in 200 Liters of water.
<b>Ginger</b>	<b>Planting</b>	Collect good quality ginger Rhizomes namely Suprava, Suruchi, Suravi. Seed rate is 6-8 q/ac. Each piece should be 20 g and contain 2-3 eyes. Treat the seeds before planting in the field.
<b>Turmeric</b>	<b>Planting</b>	Sow good varieties like Roma, Suroma, Ranga and Rashmi. Seed rate is 8-10 q/ac. The seed material should contain 2-3 eyes and of 20-25g per piece.
<b>Animal</b>		Due to rise in temperature provide the animal shed clean and sufficient drinking water. Maintain the cleanliness in the animal shed. Do not leave the animals during hot time of the day. Treat around 10 kg rhizome in 10 liters of water mixed with 40 g Mancozeb and 10 g Carbendazim and dry it under shade.
<b>Vegetables</b>	<b>Vegetative growth</b>	Due to temperature effect in brinjal Foot & Shoot borer can be managed by application of Neem based pesticides 330 ppm @ 1 liter per acre. Also due to weather there may attack of Mites, to manage this spray Dicofol @ 3 ml/liter of water.
<b>Pisciculture</b>	<b>Growth stage</b>	Due to rise in temperature, survey for oxygen deficit if any.
<b>Onion</b>	<b>Harvesting stage</b>	Harvest the onion leaving 2-3 cm from the neck and cut all the roots and dry it under shed.

For any problem related to agriculture sector, please dial the toll free number 1551.

For better agricultural benefits use, **OUAT KALINGA** Products (Seedlings, Saapplings and QPM, Tissue culture plants, Fruits and vegetables, value added products, Fingerlings/Yearlings, Poultry chicks, Breeds, Mushroom and spwn, Biofertilizer, Biopesticides, Vermi & Vermicompost and Farm implements)

**IMD weather forecast & Agrometeorological Advisory of Odisha location is now available on Meghdoot mobile app in English & Odia Language.**

Download: (Andriod: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>)

ବି : ଦ୍ରଷ୍ଟବ୍ୟ :ଓଡ଼ିଆ ଶାଖା ଚିପଲିନା ଓ ଗନ୍ଧାରିପାଲି ବିହନ ଗବେଷଣା ଫାର୍ମ ଠାରେ ସ୍ୱର୍ଣ୍ଣ ଧାନ ବିହନ ୨୦ କିଗ୍ରା ବସ୍ତା ପିଛା ଟ. ୭୧୨ ୪୦ . ପଇସା ଚାର୍ଜ ନିମିତ୍ତ ବିକ୍ରି ହେଉଅଛି ।

**Nodal Officer**