



Gramin Krishi Mausam Sewa
India Meteorological Department
Odisha University of Agriculture and Technology
Ranital, Bhadrak, - 756111



WeekNo.34

No-68(Eng.Bulletin)
Dt.-26.08.2025

District-Jajpur(North-Eastern Coastal Plain Zone)

Forecast (Upto 31.08.2025)
Given by Met. Centre,IMD,Bhubaneswar

DISTRICT: Jajpur– As per IMD forecast, the district is likely to receive **light rain for next 5 days** with generally cloudy sky.The wind will blow in a speed within 03-09km/h. The maximum temperature and minimum temperature will likely to remain 29-30°C and 23-24°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
JAJPUR	2025-08-27	8	29	23	5	91	66	3	302
	2025-08-28	8	29	24	5	91	65	9	300
	2025-08-29	7	30	24	5	91	68	6	283
	2025-08-30	7	30	23	5	91	68	7	234
	2025-08-31	7	29	23	5	92	73	6	237

For further information,contact theMet.Centre,AerodromArea,IMD,Bhubaneswar,Tel.#0674-2596116.

Crop and Livestock Specific Agromet Advisory

General Advisory	<ul style="list-style-type: none"> ➤ Farmers are advised to maintain optimum water level & monitor the incidence of Yellow Stem Borer, Leaf folder & Mealy bug in rice crop. ➤ Keep the drainage channels open to drain out excess rain water from crop fields. ➤ Considering the light rainfall forecast, farmer should take proper care during any spraying activities. ➤ The smart phone holders can also get location specific weather information from Meghdoot & Damini Apps. 	
Crop Name	Crop Stage	Agromet Advisory
Kharif Rice	Active tillering	<ul style="list-style-type: none"> ➤ To control Sheath rot in paddy, application of Hexaconazole @ 1ml/litre recommended. ➤ If the damage of stem borer is observed above ETL i.e. 5% dead heart & 10% leaf damage respectively, soil application of Rynaxypyr 0.4 G@4 kg/acre or Flubendiamide 0.7 G 5kg/acre is advised. ➤ Top dressing of fertilizer is advised 16kg nitrogen per acre within 21 days after transplanting. ➤ Strengthen the bunds to maintain the water level 3 to 5 cm in rice field.
Jute	Vegetative stage	<ul style="list-style-type: none"> ➤ Current weather is favourable for attack of insect pest like semilooper, Bihari hair caterpillar and other foliage feeders. To control these insects, spray Lamba cyhalothrin 4.9CS @ 250 ml/acre.
Sugarcane	Vegetative stage	<ul style="list-style-type: none"> ➤ Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. ➤ To control Top shot borer in sugarcane foliar spray of Fipronil 5SC @400 ml/acre is advised.
Vegetables	Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ Keep the drainage channel open to drain out excess water. ➤ To control cercospora leaf spot disease in okra, apply Potash fertilizer @ 12 kg per acre or spray Azoxystrobin 23% SC @ 1 ml per liter or (Hexaconazole 5%+Captan 70%WP) @ 2

		grams per liter or (Carbendazim 12% + Mancozeb 63% WP) @ 2 grams per liter of water. ➤To control the yellow vein mosaic virus disease in okra, remove the affected leaves and bury them & apply Potash fertilizer @500gm per gunth & Spraying of Diafenthuron 50%WP @ 2gm per liter or Dinotefuran 20%WG @ 0.5gm per liter of water in the afternoon recommended.
Mango	Planting	➤July and August are ideal time of mango planting. ➤Suitable varieties are Amrapalli, Dusseheri, Chousa, Baganpalli, Totapuri, Bombay green, Suvarnarekha, Himsagar, etc. ➤Seedling should be planted in pits at a distance of 10m X 10m. The pit size is 1m X 1m X 1m. One year old healthy grafted plant should be preferred. At the time of planting two basket compost, 80 g N, 100 g P ₂ O ₅ and 60 g K ₂ O fertilizer should be applied.
Pisciculture		➤Put plastic net around the pond dyke to prevent migration of fish due to over flowing of pond water.
Livestock		➤Give sufficient drinking water to animals. ➤Confined Cow & other domestic animal inside the shed and provide fodder and concentrated feed.

Mrs. Meera Mahanty
Jr. Agrometeorologist
AMFU, RRTTS, Ranital

Dr. Biswaranjan Nayak
ADR & NODAL OFFICER
AMFU, RRTTS, Ranital

For better agricultural benefits use, **OUAT KALINGA products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits & Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermi Compost and Farm Implements).**

Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for Weather forecast & Agromet advisory and “Damini” App for Lightning, Thunderstorm warning & Forecasting.

Download: (Android: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>), (iOS: <https://apps.apple.com/in/app/meghdoot/id1474048155>)