



Gramin Krishi Mausam Sewa
India Meteorological Department
Orissa University of Agriculture and Technology
RRTTS, Chiplima, Sambalpur



No. 64 (English Bulletin)

Week no - 32

Date: 12.08.2025

West Central Table Land Zone

District-Jharsuguda

**REALISED MAXIMUM & MINIMUM TEMPERATURE RECORDED IN THE OBSERVATORIES AND DISTRICT
 AVERAGE RAINFALL DURING THE PERIOD: 06.08.2025 to 12.08.2025**

DATES	MAX TEMP (°C)							MIN TEMP (°C)							RAINFALL (mm)						
	06	07	08	09	10	11	12	06	07	08	09	10	11	12	06	07	08	09	10	11	12
Jharsuguda	35	34	33	30	34	35	35	24	24	24	25	25	24	27	1.4	4.4	12.8	1.7	0.0	0.0	0.0

Weather forecast for next 5 days

Date	13-08-2025	14-08-2025	15-08-2025	16-08-2025	17-08-2025
Rainfall (mm)	7	6	4	6	7
Max Temperature (° C)	36	35	34	34	34
Min Temperature (° C)	25	24	24	24	24
Total cloud cover (okta)	5	5	5	5	5
Max Relative Humidity (%)	87	87	88	88	92
Min Relative Humidity (%)	60	66	60	61	61
Wind speed (km/h)	3	6	8	9	8
Wind direction (deg)	193	120	82	75	77

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

Weather Summary/Alert:

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is a chance of occurrence of light rainfall in some places along with partly to generally cloudy sky is expected during upcoming week. No large change in both Maximum and Minimum temperature is expected in upcoming 5 days. High relative humidity along with a wind speed of 3-9 km/h is predicted in south-westerly, south-easterly and north-easterly direction.

General Advisory

- Looking to the light rainfall condition in upcoming days, farmers should conduct agricultural activities like spraying, weeding, hoeing etc. with proper care at morning hours of the day under clear sky.
- Prevailing weather condition favours various diseases and sucking insects attack in both field as well as horticultural crops so its regular monitoring should be advised.
- Keep the animals in shed and keep them away from any electrical devices as thunderstorm/lightning is expected in upcoming week.
- Farmers should use Meghdoot mobile app for IMD weather forecast and Agro-meteorological advisory of Odisha in both English and Odia languages and farmers can also use Damini mobile app for thunderstorm and lightning information and Mausam app for location specific forecast and warning.

SMS Advisory

- Looking to the upcoming light rainfall events, farmers are advised to conduct any kind of agricultural activities with proper care at morning hours of the day when sky is clear.

Crop and livestock Specific Agromet Advisory

Name of the crop	Crop Stage	Agromet Advisories
Kharif Rice	Tillering	• Looking to the prevailing weather condition, farmers are advised to monitor the attack of stem borer and leaf folder in Paddy fields. To

		control these insects, keep 3 pheromone traps/acre of rice field for monitoring the stem borer and leaf folder infestations. Whenever the number of male moths/trap reaches 4 or 5, apply Azadirachtin 0.15% EC @ 800 ml/acre or, Chlorantraniliprole 0.4% GR @ 4 kg/acre mixing with sand at 1:1 ratio or, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or, Cartap hydrochloride 4G @ 10 kg/acre or, Flubendiamide 20 WG 50 g/acre in 200 litres of water.
Maize	Vegetative stage	<ul style="list-style-type: none"> Stem borer and Fall army worm attack was observed in some maize fields so farmers are advised to spray Chlorantraniliprole @ 0.4ml or Emamectin benzoate @ 0.4g per liter of water under clear weather condition to control these insects.
Mango	Vegetative growth (New orchard)	<ul style="list-style-type: none"> Due to prevailing weather condition, the attack of leaf eating caterpillar may noticed in some newly planted Mango crop, so farmers are advised to monitor their orchard regularly.
	Old orchard	<ul style="list-style-type: none"> The spider web should be cleaned and affected portion should be cut and burned. Spray Copper oxychloride @ 3g per liter of water under clear weather condition for control of red rust and anthracnose in mango.
Brinjal, Tomato	Fruiting	<ul style="list-style-type: none"> If wilting problem noticed in field, then application of Carbendazim @ 2g per liter of water followed by Hexaconazole @ 2ml per liter of water one week after first application should be advised under clear weather condition. Past weather condition is favorable for fruit borer & fruit and shoot borer attack in Tomato and Brinjal respectively so its regular monitoring should be advised. If attack observed, then spraying of Emamectin benzoate @ 0.4g or Chlorantraniliprole @ 0.3ml or Tetraniliprole @ 0.5ml per liter of water should be advised.
Chilli	Nursery management/ Planting	<ul style="list-style-type: none"> Monitoring against attack of sucking insects likes aphids, thrips etc. should be advised because prevailing weather condition is favorable for its attack.
Cows and Buffalos		<ul style="list-style-type: none"> Some precautions after milking: After milking, cows and buffalos like animals feel tired, so they want to sit down immediately, but at that time the udder is partially open, so external pathogens may easily enter the udder, which further increases the risk of Mastitis disease. Therefore, farmers should be careful about this and take proper care so that the animals do not sit on the floor immediately after milking.
Goat		<ul style="list-style-type: none"> PPR vaccine should be given to animals to protect them this virus. Weak, debilitated or infested animals should not be vaccinated. Farmers should avoid stressing vaccinated animals (e.g., transportation, inclement weather) for at least 3 weeks.
Poultry		<ul style="list-style-type: none"> To prevent the occurrence of Ranikhet disease in poultry farm, vaccinating the chicks against this disease at 1-7 days old and then again at 20-28 days old stage and also contact the nearest animal health center if necessary.

Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima	Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima
--	---