



West Central Table Land Zone

District-Bargarh

REALISED MAXIMUM & MINIMUM TEMPERATURE RECORDED IN THE OBSERVATORIES AND DISTRICT
AVERAGE RAINFALL DURING THE PERIOD: 03.09.2025 to 09.09.2025

DATES	MAX TEMP (°C)							MIN TEMP (°C)							RAINFALL (mm)						
	03	04	05	06	07	08	09	03	04	05	06	07	08	09	03	04	05	06	07	08	09
Bargarh	31	30	29	30	32	32	32	25	24	25	25	26	26	26	16	36	36	5	3	11	4

Weather forecast for next 5 days

Date	10-09-2025	11-09-2025	12-09-2025	13-09-2025	14-09-2025
Rainfall (mm)	8	7	8	8	7
Max Temperature (° C)	33	33	32	32	32
Min Temperature (° C)	26	26	25	25	25
Total cloud cover (okta)	4	5	4	5	5
Max Relative Humidity (%)	92	91	92	93	95
Min Relative Humidity (%)	63	63	66	71	66
Wind speed (km/h)	2	3	0	4	5
Wind direction (deg)	191	247	360	17	106

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

Weather Summary/Alert:

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is a chance of occurrence of light rainfall along with partly to generally cloudy sky is expected during upcoming week. No large change in both Maximum and Minimum temperature is expected in upcoming 5 days. High relative humidity along with a wind speed of 0-5 km/h is predicted in south-westerly, northerly, north-easterly and south-easterly direction.

General Advisory

- Keep the animals in safe places and don't tie them in open places during warning period.
- Farmers should use Meghdoot mobile app for IMD weather forecast and Agro-meteorological advisory of Odisha in both English and Odia languages and farmers can also use Damini mobile app for thunderstorm and lightning information and Mausam app for location specific forecast and warning.

SMS Advisory

- Monitor the crop against sucking insects, fruit fly, fruit borer, fruit and shoot borer etc. attack in vegetable crops.

Crop and livestock Specific Agromet Advisory

Name of the crop	Crop Stage	Agromet Advisories
Kharif Rice	Active Tillering / Panicle initiation	<ul style="list-style-type: none"> • If Bacterial blight / Bacterial leaf streak incidence noticed, then application of Plantomycin @ 1 g/litre along with Copper oxychloride @1 g/litre of water should be advised when sky is clear. • To monitor the attack of stem borer and leaf folder in Paddy fields, farmers should install 4-5 pheromone traps/acre. Whenever the number of male moths/trap reaches 4 or 5, apply Azadirachtin 0.15% EC @ 800 ml/acre or, Chlorantraniliprole 0.4% GR @ 4 kg/acre mixing with sand at 1:1 ratio or, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200

		litres of water or, Cartap hydrochloride 4G @ 10 kg/acre or, Flubendiamide 20 WG 50 g/acre or Tetraniliprole 200 SC (18.18% w/w) @ 100 - 120 ml/acre in 200 liters of water.
Maize	Vegetative stage	<ul style="list-style-type: none"> If stem borer and fall army worm attack was observed in maize fields then farmers should spray Chlorantraniliprole 18.5 SC @ 0.4ml or Emamectin benzoate 5% SG @ 0.4g per liter of water to control these insects under clear weather condition.
Banana	Vegetative growth (New orchard)	<ul style="list-style-type: none"> Due to past waterlogged condition in some newly planted Banana orchard, root rot disease may observed in some plants, so its regular monitoring should be advised.
Vegetables	Vegetative / Flowering/ Fruiting stage	<ul style="list-style-type: none"> Fungal disease attack may occurred in upcoming days due to prevailing weather condition so its regular monitoring should be advised. Monitor the crop against fruit fly, fruit and shoot borer, fruit borer etc. attack in vegetable crops. If sucking insects likes aphids, thrips etc. attack observed, then farmers should spray Imidacloprid 17.8 SL @ 0.3 ml or Fipronil 5 SC @ 3ml per liter of water under clear sky.
Cows and Buffalos		<ul style="list-style-type: none"> Looking to the upcoming rainfall and thunderstorm events, farmers are advised to keep the animals away from any electrical devices and keep them in safe places under shed and avoid direct entry of rain water to the shed. To prevent multiplication of mosquitoes, flies and various microorganisms in the animal shed, farmers should always keep it dry and clean and also monitor the animals against attack of FMD and Mastitis diseases.
Poultry		<ul style="list-style-type: none"> In the case of poultry, respiratory diseases, diarrhea and intestinal diseases are common in this time. Since drinking water gets contaminated during rainy days, it is necessary to ensure the provision of clean drinking water to the poultry birds. Farmers should also make provision to ensure proper ventilation in poultry farm as well as make arrangements to prevent direct entry of rain water into the shed.

Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima	Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima
--	---

- For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)
- IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information and **MAUSAM app** for location specific forecast and warning.