



**Gramin Krishi Mausam Sewa**  
**India Meteorological Department**  
**Orissa University of Agriculture and Technology**  
**RRTTS, Chiplima, Sambalpur**



No. 73 (English Bulletin)

Week no - 37

Date: 12.09.2025

**West Central Table Land Zone**

**District-Boudh**

**REALISED MAXIMUM & MINIMUM TEMPERATURE RECORDED IN THE OBSERVATORIES AND DISTRICT**  
**AVERAGE RAINFALL DURING THE PERIOD: 06.09.2025 to 12.09.2025**

DATES	MAX TEMP (°C)							MIN TEMP (°C)							RAINFALL (mm)						
	06	07	08	09	10	11	12	06	07	08	09	10	11	12	06	07	08	09	10	11	12
<b>Boudh</b>	33	33	34	34	34	34	33	24	24	24	24	24	25	25	29.8	0.0	0.0	0.0	0.0	0.0	13.0

**Weather forecast for next 5 days**

<b>Date</b>	<b>13-09-2025</b>	<b>14-09-2025</b>	<b>15-09-2025</b>	<b>16-09-2025</b>	<b>17-09-2025</b>
<b>Rainfall (mm)</b>	13	10	8	7	7
<b>Max Temperature (° C)</b>	32	33	33	32	32
<b>Min Temperature (° C)</b>	25	25	24	24	24
<b>Total cloud cover (okta)</b>	5	5	5	5	5
<b>Max Relative Humidity (%)</b>	93	94	95	95	92
<b>Min Relative Humidity (%)</b>	63	60	63	68	68
<b>Wind speed (km/h)</b>	2	4	4	3	3
<b>Wind direction (deg)</b>	127	121	186	203	214

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

**Weather Summary/Alert:**

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is a chance of occurrence of light to moderate rainfall along with generally cloudy sky is expected during upcoming week. No large change in both Maximum and Minimum temperature is expected in upcoming 5 days. High relative humidity along with a wind speed of 2-4 km/h is predicted in south-easterly and south-westerly direction.

**General Advisory**

- As per the extended range forecast issued by MC, Bhubaneswar on dated:-11.09.2025, scattered to fairly widespread rain/thundershower is very likely over districts of Odisha during week-1 (12<sup>th</sup> to 18<sup>th</sup> September, 2025). Isolated to scattered rain/thundershower is very likely over the districts of Odisha during week-2 (19<sup>th</sup> to 25<sup>th</sup> September, 2025). There is a likelihood of formation of a fresh upper air cyclonic circulation/low-pressure area over west central & adjoining northwest Bay of Bengal during first half of week-2.
- Looking to the above mentioned weather condition farmers are advised to take proper care while conducting any agricultural activities like spraying, irrigation, hoeing etc. and keep their animals in safe places during warning period. Monitor the crop against sucking insects, fruit fly, fruit borer, fruit and shoot borer etc. attack.
- If possible, use Meghdoot mobile app for IMD weather forecast and Agro-meteorological advisory of Odisha in both English and Odia languages, Damini mobile app for thunderstorm and lightning information and Mausam app for location specific forecast and warning.

**SMS Advisory**

- As light to moderate rain/thundershower very likely to occur over the district in upcoming days, farmers should wait some days for irrigation as well as spraying operation and stay safe at indoor.

**Crop and livestock Specific Agromet Advisory**

Name of the crop	Crop Stage	Agromet Advisories
------------------	------------	--------------------

<b>Kharif Rice</b>	<b>Active Tillering / Panicle initiation</b>	<ul style="list-style-type: none"> <li>• If Bacterial blight / Bacterial leaf streak incidence noticed, then application of Plantomycin @ 1 g/litre along with Copper oxychloride @1 g/litre of water should be advised when sky is clear.</li> <li>• To monitor the attack of stem borer and leaf folder in Paddy fields, farmers should install 4-5 pheromone traps/acre. Whenever the number of male moths/trap reaches 4 or 5, apply Azadirachtin 0.15% EC @ 800 ml/acre or, Chlorantraniliprole 0.4% GR @ 4 kg/acre mixing with sand at 1:1 ratio or, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or, Cartap hydrochloride 4G @ 10 kg/acre or, Flubendiamide 20 WG 50 g/acre or Tetraniliprole 200 SC (18.18% w/w) @ 100 - 120 ml/acre in 200 liters of water under clear weather condition.</li> <li>• Farmers should take proper care during spraying operation or wait some days because some amount of rainfall is expected in upcoming days.</li> </ul>
<b>Maize</b>	<b>Vegetative stage/ Tasseling</b>	<ul style="list-style-type: none"> <li>• If stem borer and fall army worm attack was observed in maize fields then farmers should spray Chlorantraniliprole 18.5 SC @ 0.4ml or Emamectin benzoate 5% SG @ 0.4g per liter of water to control these insects under clear weather condition.</li> <li>• Looking to the upcoming rainfall events, if not necessary farmers should wait some days for spraying operations.</li> </ul>
<b>Banana</b>	<b>Vegetative growth (New orchard)</b>	<ul style="list-style-type: none"> <li>• Due to past waterlogged condition in some newly planted Banana orchard, root rot disease may observed in some plants, so its regular monitoring should be advised.</li> <li>• Make provision to drain out extra water from field because some amount of rainfall is expected in upcoming days also.</li> </ul>
<b>Vegetables</b>	<b>Vegetative / Flowering/ Fruiting stage</b>	<ul style="list-style-type: none"> <li>• Fungal disease attack may occurred in upcoming days due to prevailing weather condition so its regular monitoring should be advised.</li> <li>• Monitor the crop against fruit fly, fruit and shoot borer, fruit borer etc. attack in vegetable crops.</li> <li>• If sucking insects likes aphids, thrips etc. attack observed, then farmers should spray Imidacloprid 17.8 SL @ 0.3 ml or Fipronil 5 SC @ 3ml per liter of water under clear sky.</li> </ul>
<b>Cows and Buffalos</b>		<ul style="list-style-type: none"> <li>• Looking to the upcoming rainfall and thunderstorm events, farmers are advised to keep the animals away from any electrical devices and keep them in safe places under shed and avoid direct entry of rain water to the shed.</li> <li>• To prevent multiplication of mosquitoes, flies and various microorganisms in the animal shed, farmers should always keep it dry and clean and also monitor the animals against attack of FMD and Mastitis diseases.</li> </ul>
<b>Poultry</b>		<ul style="list-style-type: none"> <li>• In the case of poultry, respiratory diseases, diarrhea and intestinal diseases are common at this time and farmers are advised to monitor the farm against Ranikhet disease. Ensure the provision of clean drinking water to the poultry birds.</li> <li>• Farmers should also make provision to ensure proper ventilation in poultry farm as well as make arrangements to prevent direct entry of rain water into the shed.</li> </ul>

<b>Ms. Jubuli Sahu</b> <b>Junior Agrometeorologist,</b> <b>GKMS, RRTTS, Chiplima</b>	<b>Dr. Sanjukta Mohapatra</b> <b>ADR cum Nodal Officer</b> <b>RRTTS, Chiplima</b>
--	---

- For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)

- IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information and **MAUSAM app** for location specific forecast and warning.