



West Central Table Land Zone

District-Bargarh

REALISED MAXIMUM & MINIMUM TEMPERATURE RECORDED IN THE OBSERVATORIES AND DISTRICT
AVERAGE RAINFALL DURING THE PERIOD: 25.10.2025 to 31.10.2025

DATES	MAX TEMP (°C)							MIN TEMP (°C)							RAINFALL (mm)						
	25	26	27	28	29	30	31	25	26	27	28	29	30	31	25	26	27	28	29	30	31
Bargarh	34	32	33	32	32	32	32	25	25	25	24	24	24	25	0	0	0	5	4	2	1

Weather forecast for next 5 days

Date	01-11-2025	02-11-2025	03-11-2025	04-11-2025	05-11-2025
Rainfall (mm)	5	3	2	1	1
Max Temperature (° C)	32	33	33	33	34
Min Temperature (° C)	23	23	23	22	22
Total cloud cover (okta)	4	3	3	2	2
Max Relative Humidity (%)	91	91	92	91	85
Min Relative Humidity (%)	62	61	63	59	55
Wind speed (km/h)	6	4	5	4	3
Wind direction (deg)	220	259	299	355	13

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

Weather Summary/Alert:

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is less chance of occurrence of rainfall but very light to light rain may occur at some places during upcoming week and partly clear to partly cloudy sky is expected during that period. No large change in both Maximum and Minimum temperature is expected in upcoming 5 days. High relative humidity along with a wind speed of 3-6 km/h is predicted in south-westerly, north-westerly and north-easterly direction.

General Advisory

- The excess water should be drain out immediately from the field having standing crops and not ready to harvest.
- Due to heavy rainfall the incidence of disease pest may increase. Regularly visit the field to know the incidence of diseases and insects pest. Accordingly take need based pest management measure immediately.
- Farmers those are using android mobile should use Meghdoot mobile app for IMD weather forecast and Agro-meteorological advisory of Odisha in both English and Odia languages, Damini mobile app for thunderstorm and lightning information and Mausam app for location specific forecast and warning.

SMS Advisory

- Farmers should start sowing of rabi non-paddy crop by taking the advantage of past rainfall condition.

Crop and livestock Specific Agromet Advisory

Name of the crop	Crop Stage	Agromet Advisories
Kharif Rice	Grain filling/ Dough/ Maturity/ Harvesting	<ul style="list-style-type: none"> • Due to prevailing weather condition, there may be chances of infestation of Brown Plant hopper (BPH), Gundhi bug in long duration varieties of rice or very late planted rice and ear cutting caterpillar in matured/harvested crop in the field so its regular monitoring should be advised. • Harvesting of Paddy crop should be advised when 80-85% of the grains are matured. Harvesting should be done either manually by sickle or by

		<p>using combine harvester or reaper.</p> <ul style="list-style-type: none"> After harvesting grains need to be sun-dried to 14% moisture content for consumption purpose and for seed purpose it should be dried to 12% moisture for better shelf-life.
Rapeseed/ Mustard	Variety selection/ Sowing	<ul style="list-style-type: none"> Choose suitable varieties like Pusa bold, Binaya, Pusa Bahar, Baruna, Pusa Jaikisan, JD-6. Toria- M 29, Anuradha, Parbati etc. should be advised for sowing under proper moisture condition. Make provision to drain out excess water from already sown crop fields if any.
Groundnut	Variety selection/ Sowing	<ul style="list-style-type: none"> Farmers should use improved varieties of Groundnut such as T.A.G. -24, T.G.-3, T.G.-38, Kadri-5, 6, 9, C.O.-7, Devi etc. 100 to 130 days old varieties. Ensure seed treatment before sowing and treat with bacteria or phospho-culture 7 days after treatment. 40 to 50 kg of seed or 60 to 70 kg of pod is required per acre. The seeds should be sown within 2 to 3 days of decortication.
Mango	Orchard Management	<ul style="list-style-type: none"> Ensure good drainage facilities in mango orchard. Apply recommended fertilizer dose around the tree's trunk (from 45-90 cm away, out to the drip line) and incorporate it into the soil. Pruning of any dead or diseased branches should be advised and then apply copper oxychloride @ 3g per liter of water twice at a 15-day interval after pruning to prevent further infections.
Tomato	Nursery bed preparation/ Seedling/ Planting	<ul style="list-style-type: none"> Prepare a nursery bed of 3m length, 1m breadth and 15-20 cm height and apply 15-30 kg well decomposed FYM and use 150 grams of chlorpyrifos dust to each nursery bed seven days before sowing. Monitor the seedlings against various insects and diseases and also planting of healthy seedlings should be done with a spacing of 60 cm × 45 cm in tomato field.
Cows and Buffalos		<ul style="list-style-type: none"> Looking to the upcoming thunderstorm events, farmers are advised to keep the animals away from any electrical devices and keep them in safe places under shed and avoid direct entry of rain water to the shed and also monitor the animals against attack of FMD and Mastitis diseases.

Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima	Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima
--	---

- For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)
- IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information and **MAUSAM app** for location specific forecast and warning.