



**Gramin Krishi Mausam Sewa**  
**India Meteorological Department**  
**Odisha University of Agriculture and Technology**  
**Ranital, Bhadrak, - 756111**



WeekNo.46

No-91(Eng.Bulletin)  
Dt.-14.11.2025

District-Bhadrak (North-Eastern Coastal Plain Zone)

**Weather condition during last week at Bhadrak (08.11.2025--14.11.2025)**

Rainfall (mm)	0.0
Maximum temperature (°C)	29.0-31.0
Minimum temperature (°C)	17.0-19.0

Forecast (Upto 19.11.2025)  
Given by Met.Centre,IMD,Bhubaneswar

**DISTRICT: Bhadrak** As per IMD forecast, the district is likely to receive **dry weather for next 5 days** with clear sky. The wind will blow in a speed within 09-11km/h. The maximum temperature and minimum temperature will likely to remain 29-30°C and 17-19°C to respectively for the next five days.

DISTRICT	Date	Rainfall (mm)	T-MAX (C)	T-MIN (C)	Cloud Cover	Rh Max (%)	Rh Min (%)	Wind speed (kmph)	Wind Direction (deg)
BHADRAK	2025-11-15	0	29	17	1	64	32	10	351
	2025-11-16	0	29	17	1	62	34	9	356
	2025-11-17	0	30	18	1	61	34	11	2
	2025-11-18	0	30	18	2	63	40	11	2
	2025-11-19	0	30	19	2	74	48	11	344

For further information, contact the Met.Centre,aerodrome Area,IMD,Bhubaneswar,Tel.#0674-2596116.

**Crop and Livestock Specific Agromet Advisory**

<b>General Advisory</b>		<ul style="list-style-type: none"> <li>➤ Monitor the activity of Gundy bug in rice field and install rotten snail trap for mass trapping and killing of the pest.</li> <li>➤ Provide irrigation at the critical stages of growth of standing rice &amp; other vegetable crops.</li> <li>➤ Taking the advantage of store soil moisture, farmer should start land preparation for rabi crops like mustard, toria, sunflower and sowing should be completed soon to get optimum yield.</li> <li>➤ Visit to field regularly to observe the infestation levels of Stem borer, Leaf folder, Gundy bug and the incidence of Blast.</li> </ul>
<b>Crop Name</b>	<b>Crop Stage</b>	<b>Agromet Advisory</b>
<b>Kharif Rice</b>	Booting/ Flowering	<ul style="list-style-type: none"> <li>➤ Prevailing warm and humid weather may increase the incidence of Blast disease in rice so apply (Tebuconazole + Trifloxystrobin) @100g/acre along with plantomycin@200g/acre recommended.</li> <li>➤ Prevailing microclimate condition may increase the population of Plant Hoppers. If the BPH &amp; WBPH infestation is at initial stage alternating drying &amp; wetting of field &amp; foliar application of neem based formulation 1500ppm (600ml /acre), Flonicamid@80g/acre or Dinotefuron@80g/acre recommended. If the population is around ETL(8-10)/hill spraying of (Ethiprol + pymetrozine)@170g/acre recommended.</li> <li>➤ Monitor the population level of WBPH &amp; BPH by regular field visit.</li> </ul>
<b>Rapeseed and Mustard</b>	Land preparation	<ul style="list-style-type: none"> <li>➤ Taking the advantage of store soil moisture, land preparation and sowing of the crop should be completed soon to get optimum yield.</li> <li>➤ Suitable varieties: Toria- Anuradha, Parbati, Sushree, jayanti</li> <li>➤ Indian mustard- NRCHB 101, Sampurna</li> </ul>

		➤Seed rate- 8-10 Kg/ha, Spacing- 30 cm X 10 cm
<b>Vegetables</b>	Nursery /Flowering/ Fruiting	<p>➤Select suitable varieties of Cole crops, Chilli and Brinjal and sow the seeds on the well pulverized nursery bed. Fully decomposed FYM or compost should be added to the nursery bed before sowing.</p> <p>➤For minimizing the infestation of sucking insect pests like white fly, leaf hopper thrips and aphid in vegetables nursery, spray Thiamethoxam @ 0.4g/litre of water.</p> <p>➤If the seedlings of early duration varieties of vegetables are damaged , treat the seedlings with (Metalaxyl + Mancozeb ) @ 2g/li of water and Plantomycin @ 1g/liter of water.</p>
<b>Fruits</b>	Flowering/ Fruiting	❖ <b>Coconut</b> -Monitor the activity of the Red palm weevil and Rhinoceros beetle by placing pheromone trap. Spray Lambda cyhalothrin @ 1.2ml/lit of water to the nearby manure pits for killing the developing stages of Rhinoceros beetle.
<b>Livestock</b>		<p>➤If the animals have not yet been vaccinated against FMD, Haemorrhagic Septicaemia, Black Quarter, Enterotoxemia etc., these must be done.</p> <p>➤Keep shed clean and dry to protect animals from disease and insects</p>

Mrs. Meera Mahanty  
Jr. Agrometeorologist  
AMFU, RRTTS, Ranital

Dr.Biswaranjan Nayak  
ADR & NODAL OFFICER  
AMFU,RRTTS, Ranital

**For better agricultural benefits use, **OUAT KALINGA** products (Seedlings, Saplings & QPM, Tissue Culture Plants, Fruits &Vegetables, Value Added Products, Fingerlings/ Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides,Vermi&Vermi Compost and Farm Implements).**

Kindly download and use the “MAUSAM” App for location-specific forecast and warning, “Meghdoot” App for Weather forecast &Agromet advisory and “Damini” App for Lightning ,Thunderstorm warning & Forecasting.

**Download:(Android:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>),(iOS:<https://apps.apple.com/in/app/meghdoot/id1474048155>)**