



Gramin Krishi Mausam Sewa
India Meteorological Department
Orissa University of Agriculture and Technology
RRTTS, Chiplima, Sambalpur



No. 94 (English Bulletin)

Week no - 47

Date: 25.11.2025

West Central Table Land Zone

District-Boudh

REALISED MAXIMUM & MINIMUM TEMPERATURE RECORDED IN THE OBSERVATORIES AND DISTRICT
 AVERAGE RAINFALL DURING THE PERIOD: 19.11.2025 to 25.11.2025

DATES	MAX TEMP (°C)							MIN TEMP (°C)							RAINFALL (mm)						
	19	20	21	22	23	24	25	19	20	21	22	23	24	25	19	20	21	22	23	24	25
Boudh	27	27	27	27	27	27	27	18	18	18	18	18	-	18	0	0	0	0	0	0	0

Weather forecast for next 5 days

Date	26-11-2025	27-11-2025	28-11-2025	29-11-2025	30-11-2025
Rainfall (mm)	0	0	0	0	0
Max Temperature (° C)	26	25	26	27	28
Min Temperature (° C)	18	17	16	17	18
Total cloud cover (okta)	0	0	2	1	1
Max Relative Humidity (%)	90	86	71	76	82
Min Relative Humidity (%)	39	32	26	33	38
Wind speed (km/h)	2	0	1	2	3
Wind direction (deg)	154	360	243	243	225

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

Weather Summary/Alert:

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is less chance of occurrence of rainfall during upcoming week and clear to mainly clear sky is expected during that period. No large change in both Maximum and Minimum temperature is expected in upcoming 5 days. Decrease in relative humidity along with a wind speed of 0-3 km/his predicted in south-south-easterly, north, west-south-westerly and south-westerly direction.

General Advisory

- Looking to the upcoming dry weather condition, harvesting of matured paddy and non-paddy crops should be advised as soon as possible. Farmers should maintain adequate moisture in standing crop (vegetative to reproductive stage) fields.
- To protect the crops from cold, farmers should apply light and frequent irrigation / sprinkler irrigation in the evening hours.
- Farmers those are using android mobile should use Meghdoot mobile app for IMD weather forecast and agro-meteorological advisory of Odisha in both Odia languages. Damini mobile app for thunderstorm and lightning information and Mausam app for location specific forecast and warning.

SMS Advisory

- Do not disturb the soil; loose surface reduces the conduction of heat from lower surface.

Cropand livestock Specific Agromet Advisory

Name of the crop	Crop Stage	Agromet Advisories
Kharif Rice	Maturity/ Harvesting	<ul style="list-style-type: none"> Harvesting of Paddy crop should be advised when 80-85% of the grains are matured. Harvesting should be done either manually by sickle or by using combine harvester or reaper. For safe storage of paddy/rice, use ‘Super Grain Bag’ which is helpful for retaining the quality, texture, color, aroma and taste for longer period of time and also prevents insect pest infestation.

		<ul style="list-style-type: none"> • Soon after noticing the infestation of the stored grain insect pest, take up fumigation by using Aluminium phosphide (do not use in dwelling houses) tablets @ of 3 tablets/ ton grain (total 9 gm of tablets) in fairly air tight containers or by covering with thick tarpaulin leaving no gaps. The tablets should be wrapped in cotton pouches before placing them in the stacks. All the corners of plastic cover should be plastered with 6-inch-thick layer of mud/ adhesive tapes to prevent leakage of gas. Minimum exposure period is for about 7-10 days.
Rapeseed/ Mustard	Variety selection/ Sowing	<ul style="list-style-type: none"> • Choose suitable varieties like Pusa bold, Binaya, Pusa Bahar, Baruna, Pusa Jaikisan, JD-6. Toria- M 29, Anuradha, Parbati etc. should be advised for sowing under proper moisture condition. • 3 to 4 kg of seeds is required per acre. For good yield, if possible, seed should be sown in rows and spacing should be a distance of 30 to 45 cm between rows and 10 cm between plants.
Groundnut	Variety selection/ Sowing	<ul style="list-style-type: none"> • Farmers should use improved varieties of Groundnut such as T.A.G. -24, T.G.-3, T.G.-38, Kadri-5,6,9, C.O.-7, Devi etc. 100 to 130 days old varieties. • Ensure seed treatment before sowing and treat with bacteria or phospho-culture 7 days after treatment. 40 to 50 kg of seed or 60 to 70 kg of pod is required per acre. The seeds should be sown within 2 to 3 days of decortications.
Cole crops	Nursery bed preparation / Sowing	<ul style="list-style-type: none"> • Prepare raised nursery beds (about 10-15 cm high, 1 m wide, and of a convenient length) to facilitate drainage and intercultural operations and applied well-decomposed farmyard manure (FYM) or compost at a rate of 2-3 kg/m². • Drench the beds with any suitable fungicide solution to prevent damping-off disease and hot water treatment at 50°C for 30 minutes is also effective against black rot. • Sow seeds thinly in shallow furrows (1-2 cm deep) with a spacing of 5-7 cm between lines. Cover the seeds with a fine layer of soil and provide light watering using water can. • Cover the beds with dry straw or grass until germination is complete to maintain moisture and temperature, removing the cover immediately after seedling emergence.
Cows and Buffalos		<ul style="list-style-type: none"> • Keeping in view of decrease in minimum temperature in upcoming days, animal should kept inside the sheds during night and provide dry bedding to protect them from cold. • Mineral mixture along with salt should be given regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.
Poultry		<ul style="list-style-type: none"> • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

Ms. Jubuli Sahu Junior Agrometeorologist, GKMS, RRTTS, Chiplima	Dr. Sanjukta Mohapatra ADR cum Nodal Officer RRTTS, Chiplima
--	---

- For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)
- IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use

DAMINI mobile app(<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information and **MAUSAM app** for location specific forecast and warning.