



**West Central Table Land Zone**

**District-Jharsuguda**  
Weather forecast for next 5 days

Date	07-01-2026	08-01-2026	09-01-2026	10-01-2026	11-01-2026
Rainfall (mm)	0	0	0	0	0
Max Temperature (° C)	26	27	27	28	28
Min Temperature (° C)	5	5	6	6	7
Total cloud cover (okta)	0	1	2	4	3
Max Relative Humidity (%)	56	55	53	56	59
Min Relative Humidity (%)	21	22	22	21	26
Wind speed (km/h)	8	8	10	9	4
Wind direction (deg)	41	40	46	34	73

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

**Weather Summary/Alert:**

According to medium range weather forecast of IMD, MC, Bhubaneswar, there is very less chance of occurrence of rainfall and mainly to partly cloudy sky is expected during upcoming week. No large change in Maximum temperature but fall in Minimum temperature is expected in upcoming 5 days. Moderate relative humidity along with a wind speed of 4-10 km/h is predicted in north-easterly direction.

**General Advisory**

- Looking to the prevailing cold weather condition, farmers are advised to apply light, protective irrigation to the crops to protect them from cold.
- Past cold and foggy conditions may lead to increased infestation of certain pests, such as aphids on vegetables, beans, and mustard. High humidity and persistent leaf wetness create a conducive environment for fungal and other diseases. So, its regular monitoring should be advised.
- Where irrigation facilities are available, farmers should complete sowing of short duration rabi crops like Mustard, Chickpea, Groundnut Sesame, Green gram/ Black gram, Cowpea etc. in medium/shallow lowlands utilizing the available soil moisture in the field and also start nursery sowing of Summer Paddy as soon as possible.

**SMS Advisory**

- Farmers are advised not to burn the Paddy straw in fields.

**Crop and livestock Specific Agromet Advisory**

Name of the crop	Crop Stage	Agromet Advisories
<b>Rabi Rice</b>	<b>Nursery bed preparation/ sowing</b>	<ul style="list-style-type: none"> <li>Nursery seed beds should be prepared in puddled soil of 120 cm wide, 10 cm height and of convenient length with 30 cm gap in between two beds. About 400 m<sup>2</sup> nursery area is required for transplanting one-acre main field.</li> <li>Before sowing, seed treatment should be done with Carbendazim 50 WP @ 1.5 g/kg of seed or Trichoderma formulation @ 10 g of /kg of seed.</li> <li>Looking to the prevailing cold weather condition, farmers are advised to cover the nursery beds with polythene sheet during night in rice nursery and remove in the morning. Irrigate the nursery beds in the evening and drain out the water in the morning.</li> </ul>
<b>Ground nut</b>	<b>Sowing/ Vegetative</b>	<ul style="list-style-type: none"> <li>Past weather condition (cloudy and foggy) is favourable for sucking insect attack in groundnut crop so its regular monitoring should be</li> </ul>

	<b>stage</b>	<p>advised. If attack observed, then at initial stage spraying of Azadiractin or neem-based insecticide 1500 ppm @ 600 ml in 200 liters of water per acre.</p> <ul style="list-style-type: none"> <li>For chemical management, apply 40 grams of thiamethoxam or 80 grams of Emamectin benzoate per acre in 200 liters of water.</li> </ul>
<b>Cole crop</b>	<b>Vegetative Stage/ Fruiting</b>	<ul style="list-style-type: none"> <li>Straw can be used as mulching material in vegetable crops to soil maintain moisture as well as soil temperature in field.</li> <li>Regular monitoring against the attack of Diamond back moth in cole crops should be advised. If attack observed then spraying of neem oil @ 5 ml or Emamectin Benzoate @ 0.4 g per liter water should be advised.</li> <li>Harvesting of matured crops as well as one light irrigation in crops which are in vegetative growth stage should be advised because dry weather is expected in upcoming week.</li> </ul>
<b>Chilli</b>	<b>Vegetative stage</b>	<p><b>Leaf curl disease:</b></p> <ul style="list-style-type: none"> <li><b>Symptoms:</b> The plant does not grow well due to this disease. When the leaves are affected by the Thrips, the leaves curl upwards and these insects are present on the upper part of the young leaves and when the Aphid attacks, the leaves curl downwards.</li> <li><b>Management:</b> For its management, blue sticky traps are used to attract the Thrips. Apart from this, initially spray 5 ml of neem based insecticide mixed in one liter of water. If still persist, then spraying of chemical insecticides like Spinosad @ 4 ml in 10 liters of water or Fipronil 5% SC @ 2 ml per liter of water at an interval of 7-10 days should be advised.</li> <li>Spraying should not be done during foggy weather.</li> </ul>
<b>Cows and Buffalos</b>		<ul style="list-style-type: none"> <li>To protect the animal from cold, they should kept inside the sheds during night and provide dry bedding (If possible, use jute bag to cover the animal and straw as bedding material).</li> <li>Increase protein level and minerals in the feed concentrate to keep the animals healthy to cope up with the cold condition. Provide mineral mixture along with salt regularly and also if possible, then use wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.</li> <li>Vaccination and deworming should be done to control different harmful diseases and stomach worms.</li> </ul>
<b>Poultry</b>		<ul style="list-style-type: none"> <li>In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.</li> <li>The drinking water should be maintained at suitable temperature to promote water intake.</li> </ul>

<p><b>Mrs. Jubuli Sahu</b>  <b>Junior Agrometeorologist,</b>  <b>GKMS, RRTTS, Chiplima</b></p>	<p><b>Dr. Sanjukta Mohapatra</b>  <b>ADR cum Nodal Officer</b>  <b>RRTTS, Chiplima</b></p>
--	--

- For better agricultural benefits, use **OUAT KALINGA** products (Seedling, Sapling & QPM, Tissue Culture Plants, Fruits and Vegetables, Value Added Products, Fingerling/Yearlings, Poultry chicks, Breeds, Mushroom & Spawn, Bio-Fertilizer, Bio-Pesticides, Vermi & Vermicompost and Farm Implements.)
- IMD Weather forecast and Agro-meteorological advisory of Odisha location is now available on **MEGHDOOT mobile app** (<https://play.google.com/store/apps/details?id=com.aas.meghdoot>) in both English and Odia languages and farmers can also use **DAMINI mobile app** (<https://play.google.com/store/apps/details?id=com.lightening.live.damini>) for thunderstorm and lightning information and **MAUSAM app** for location specific forecast and warning.